

Bald eagle, glaucous-winged gull, Bonaparte gull and mew gulls all fed during high wind. The Bonaparte gull was most adept.

The salt water sprayed shorelines and beyond which turned to ice.

Nov 28, 1993


Temp. 24°F. Loss of red berries of madrona dependent on velocity of wind.

2:00 P.M. waves below whitecaps.

On west side of causeway between S Fisherman Bay Lagoon and San Juan channel 11 buffleheads, 3 glaucous-winged gulls. On E side causeway; approx 300 western sandpeeps resting in sun, 2 groups of approx 20 each of northern pintail, 23 green-winged teal, 18 glaucous-wing gulls, 2 great blue herons, 6 American crows. The south end of S Fisherman Bay Lagoon in windrows of silt, ^{brownish} from main Fisherman Bay. Half lagoon in ice.

All lagoon inspected are 1/2 in ice.

no yellow color on Blakely,



Nov. 29, 1993

To Anacortes and then Chris' for Thanksgiving. Left Lopez Island via ferry 7:50 A.M. First birds in Thatcher Pass; 21 Bonaparte gulls. E Thatcher Pass 85 Bonaparte gulls, 11 glaucous-winged gulls, 8 double-crested cormorant all generally distributed, no glaucous-winged gulls on Louter Island. 1 mew gull approaching Anacortes ferry terminal. Arrive.

Checked Marsh Point. West side, no birds. Fresh water drainage in ice, North end marsh point, 56 buffleheads,