

on face of y mt and in upper rock canyon proper.
 Discovered probable hibernating log of a bear on mountain south of Eastmans cabin in Rock Canyon. A large hollow aspen had been partly covered by a rock slide and offered an excellent chamber for a retreat. At back of this old log found dried leaves and grasses, ^{and} were probably used for a bed. Reinwald also relates of an instance where a bobcat walked into a herd of sheep in brood daylight without starting any commotion among the sheep. At Wolf Cr. divide. Also a bear had been reported at Wolf Cr. as killing 15 sheep in one night.

3/16/39

Observed the first sea gull flying over Bravo City since winter months.

3/18/39

Harry Chandler relates the following observations concerning the northern migration of the red-wing blackbird. These birds I had observed several times but did not have a complete picture. Harry is located at 9th west and 4th north which was just east of the main flights. The migration lane did not conform to water channels or any apparent topographical expression of the country. "noise of migrating birds awakened me about 6:40 A.M. on Monday, Mar 13th, while waiting for fire to warm room, observed from my window the continuous stream of these northerly migrating birds. Successive flocks of about 300 birds seemed to constitute the manner of flight. Should estimate that possibly 30,000 birds passed by during the short time they were under observation. Migration stopped at about 7:10 A.M. Main migration occurred Monday followed by a lesser one on Tues. On Wednesday the migration subsided. The occurrences of these birds may have started previous to the first date of observation. Migration on Monday lasted for about 30 minutes, trending in a northerly direction. These birds were flying about 120' high." Harry also relates that he saw 3 morning clock butterflies on canal below President Harris' at Upper University Campus. He felt that the insect life is now as abundant as it will ever be. He knows of instances where the morning clock butterflies have spent the winter unaffected by the season. These were found inactive in damp cellars.