

but was certainly the eagle nest as ⁴⁰⁰³²⁵⁻²⁹ judged by the size. Other interesting observations made were: The large roosting or better, resting rocks on top of the ridge which was covered with long streaks of white wash. Several other definite resting and observation points were seen. In many sections where the sheep could not get were grasses unaffected and enjoying a rank growth. Where the grasses were burned the Oreohelot were more in evidence and found in the more protected portions of the cliffs. Discovered the only likely evidence of the nestoma on the island. It was a typical accumulation of rocks and sticks & nest material placed in a honeycomb pocket about 1/2 sq foot. This nest, ^{was} hardly large enough in size or material to accommodate a Nestoma & have wondered whether it could have possibly been some other form. The other place was a hole under rock with sticks on ground at entrance but not definitely filled in a pile. The hole had a nestoma odor. Many other caves & holes showed signs of large animal habitation as the marmot or skunk. Their droppings were generally found among all ledges. Bonneville level on these cliff outcroppings ~~are~~ is not distinct. Lower levels show rock polishing & some find beautiful sheet & gneiss polished rocks. A spring issues from the canyon just east & south of this cliff mass. From the top of this ridge took a picture (2-3-25-40) of the shore line & mountain side to the north, showing the appearance of the shore line and terminal island where cormorants were found to have nested last year. Also the Bonneville terraces are beautifully shown on end island. It is in this N.W. section of the island where the bufflo bird is generally found to occur. From near where this picture was taken collected a Hemiptera (3-3-25-40) from the surface of a cliff in a sunny exposure. A sparrow hawk flew by here. There appears to be a congregation of rovers on high point of main ridge to the east. No rattlesnakes appeared. A cast snake skin was found however but was definitely not a rattler. Dad told me he found the patchnose last year on peninsula in sage bush on east side of the island. From the west side



2-3-25-40

generally found to occur. From near where this picture was taken collected a Hemiptera (3-3-25-40) from the surface of a cliff in a sunny exposure. A sparrow hawk flew by here. There appears to be a congregation of rovers on high point of main ridge to the east. No rattlesnakes appeared. A cast snake skin was found however but was definitely not a rattler. Dad told me he found the patchnose last year on peninsula in sage bush on east side of the island. From the west side