

Rock Island, Utah Lake. Dr Beck trapped a few mammals on rock island. He remained on island that night & next day and found that the greatest activity of the gull occurred during the night with a cessation of movement during the hot part of the day. The Canadian Goose had deserted its nest. Eggs had been deserted since beginning as embryos had just begun. His mammals were handed over to me for measurement and they are as follows. and include *Peromyscus* and *Mus*.

1-5-18-40	<i>Peromyscus</i> m. son.	162-70-20.5-22grs	♂	testis 8 m. m.
2-5-18-40	<i>Peromyscus</i> m. son.	160-71-19.5-19grs	♂	testis 7 m. m.
3-5-18-40	<i>Peromyscus</i> m. son.	165-75-20-19grs	♀	
4-5-18-40	<i>Peromyscus</i> m. son.	163-72-20.1-23grs	♂	testis 7 m. m.
5-5-18-40	<i>Peromyscus</i> m. son.	166-75-20-22grs	♂	testis 8 m. m.
6-5-18-40	<i>Peromyscus</i> m. son.	147-65-19.5-20grs	♂	testis 7 m. m.
7-5-18-40	<i>Peromyscus</i> m. son.	162-73-21.4-19grs	♂	
8-5-18-40	<i>Peromyscus</i> m. son.	156-74-20.1-15grs	♂	testis 3 m. m.
9-5-18-40	<i>Peromyscus</i> m. son.	160-72-20.2	♂	testis 7 m. m.
10-5-18-40	<i>Peromyscus</i> m. son.	152-71-20	♀	
11-5-18-40	<i>Mus musculus</i>	175-91-19.5-17grs	♂	
12-5-18-40	<i>Mus musculus</i>	154-78-18-12grs.		