

of alpine <sup>from Baseelder</sup> Canyon south for 2 or 3 canyons. Mr Johnson of American Fork and Treganza made an attempt at one time. One Clark Crowe led them directly to the nest. (wonder question this last statement.) In conclusion can say that birds and mammals are not active during these spring snow storms in windy ridges but are active immediately after storms change. *Dendroica obscura* *obscura* plumage appeared to be, at least externally, water soaked.

[see page 410609-80 for June 9, 1941]

6/22/41

Beddulphe Cabin at Aspen Grove. *Corynorhynchus r. pallidus* on *Abies concolor* with 2 young. Adult covers young with partially extended wings as many birds are prone to do. Spiders over H<sub>2</sub>O catching insects treading in circles on surface. Hummer, chippy in evidence after sun had set

6/25/41

Ascent Baseelder Peak (Thermometer mt on Sleigh Runner.) Daybreak 3:50 A.M. at Provo. Robins and a Pewee or some type of flycatcher that sings most every hour of the early morning and late evening. Left Provo at 4:00 A.M. Calif Gull flying over field at 4:00 P.M. About 1/2 mile down canyon from the Alpine Power house observed a deer among the open field in flats, also a cottontail rabbit. Left the Power house at 4:36 A.M. and trended east to mouth of Phelps Canyon. Enroute to base of mt at Phelps Canyon observed the following.

- *Hedymeles melanocephalus papago* 3
- *Passerina amoena* 3
- *Pipilo maculatus montanus* 6
- *Penthestes* 1
- *Aphelocoma californica woodhousei* 1
- *Vermivora celata* (?) 3
- *Salasphorus platycercus platycercus*
- *Picus picus hudsonicus* 3
- Several crickets calling.

Cool down canyon breeze. Few isolated cloud in sky.