Melospiza melodia morphna. as usual.
Sturnella neglecta. Few during day. Lack of birds in city district but found in outer limits.
Sialia mexicana. Several.
Junco oregonus oregonus
Junco oregonus shufeldti.
Pipilo maculatus oregonus 4 or 5
Lopia curvirostra bendirei. Few in residential
Spinus pinus pinus.
Carpodacus mexicanus frontalis 4 in one group.
Dendroica audubonii audubonii. The more brilliant males and spring plumage in evidence.
Sitta carolinensis.
Thryomanes bewicki calophonus.
Zonotrichia leucophrys pugetensis. 2 birds
Psaltriparus minimus minimus.
Colaptes cafer cafer groups of 1-1-1-1-1-1-1
Cyanocitta stelleri stelleri. 4 together
Hesperiphona vespertina brooksi. 18
Zenaidura macroura marginella 3
Anas platyrhynchos platyrhynchos 13-16 and other groups
Marecca americana 78-823 and small groups
Nettion carolinense 23-69 and other small groups.
Oxyechus vociferus vociferus. 1-1-1-1-1-1-1-1-
Corvus brachyrhynchos brachyrhynchos 2-1-2-1-1-2
Aix sponsa 1-5
Euphagus cyanocephalus 21
Accipiter cooperi 1
Penthestes atricapillus occidentalis
Melospharus rufus 11
Fulica americana 14-6
Agelaius phoeniceus 30
Ardea herodias fannini.
Anthus spinoletta
Larus glaucescens Group of about 6,000 gulls with mixed species
Oxyabatis pubescens gardneri. 1
Passerculus s. alaudinus 1
Spatula clypeata 2-2-2-4
Chaulelasmus streperus 3
Dafila acuta tzitzihoa. 4
Tyto alba 1 dead