

Setta Canadensis - usual.

Puranga ludroceana. well represented.

Empidonax traillii brewsteri. Decided decrease from 2 weeks ago.

Colaptes cafer cafer. Approx. 12 birds

Oxyechus vociferans vociferans. Decided increase particularly in the mud flats along lake shore. at south end of lake. Approx. 130 birds. Killdeers are associated with nearly all the other shore birds.

Junco oregonus shufeldti. Few.

Sporus pinus pinus.

Phasianus colubinus torquatus. Skulked several birds at 3 feet in tall grasses where they had been bedded for the night.

Euphagus cyanocephalus. Few flocks of 7-8

Butorides versicolor anythoni. Six birds widely distributed.

Carpodacus mexicanus frontalis. Increase

Zonotrichia leucophrys pugetensis.

Sporzella passerina arizonae. Approx 6

Bonhycella cedrorum

Dryobates pubescens gairdneri. Approx 8

Dendroica auduboni. Approx. 7 birds

Vireo solitarius cassinii. 2 birds.

Zenaidura m. macroura 3 birds

Vireo huttoni 1

Megascops alpestris calurus

Anas p. platyrhynchos. Approx 180

Passerculus sandwichensis. 8

Ait sponsa. Approx. 28

Empidonax difficilis difficilis (?)

Capella delicata 1

(Extremely questionable, probably Pisobia melanostris)

Tringa solitaria canadensis. (?) Fed with 2

killdeer at point where Levee Creek enters bay lake.

Bird has characteristic high arc wing beat.

Dendroica aestiva 1

Larus delawarensis. dominant. Gull flight from east and south at sundown.

Corvus b. brachyrhynchos. Increase.