

Sitta canadensis. Heard 2 during 420920-108

Regulus satrapa olivaceus. One group of this <sup>early morning</sup> form early morning among conifers; otherwise poorly represented.

Thryomanes bewickii calaptonus. Approx. 8 birds which is a better representation than former listings. One bird sang to the rising sun in a short melodious and musical manner.

Geothlypis trichas occidentalis. Three birds found in the following situations. One among typical creek bottom groups of willow, stinging nettle, blackberry vines etc with creek supplied. no grasses or cattails associated. Another bird among brush on semi-dry side hill 300' from water. Another bird in old willow pile 500' from water bordering a willow line succession along south shore line. These birds are found in environs somewhat disassociated from the swamp environ of Utah. Have often wondered if possibly it could be separated upon a physiological basis.

Spinus pinus pinus. One flock of about 20 birds, otherwise occasionally recorded in small groups of 2 or 3.

Sturnella neglecta. 2 birds only. These heard singing in early morning. Why they are not represented in a more respectable manner is beyond my knowledge.

Phasianus colchicus torquatus. 12 birds in small groups.

Junco oregonus shufeldti.

Corthylio calendula grinnelli. 1 bird in mixed flock.

Penthestes atricapillus occidentalis.

Piranga ludoviciana. Voice record only of one bird as compared to approx 20 of last week.

Zonotrichia coronata. One bird. Doubtful

Passer domesticus

Tiro solitarius cassini 4 birds

Spizella passerina arizonae. 2

Loxia curvirostra hendirei Questionable

Psaltiriparus minimus minimus. One group of 24

Oryzobates pubescens gardneri 10

Butorides virescens anthonyi. 2 only.