

exposures. Found the mole to penetrate ^{bare} ~~moist~~ flats to certain maximum zones of moisture then end abruptly. These underground chambers were not supplied with any vegetation root systems whatsoever, being extended 20-30 feet beyond the peripheral margin of growth.

10/4/42

This afternoon dropped over to Severeth Creek at end of Q street. Itinerary included creek bottom, hillside and timbered land immediately adjacent creek. Observed the following. Day clear. Occasional slight wind. Leaves falling from many trees, other leaves turning.

Cyanocitta stelleri stelleri. 12. One bird beating willows as if mad or insane

Penthestis atricapillus occidentalis.

Regulus satrapa olivaceus

Phasianus calchurus torquatus

Corythoichthys calendula gunnelli. Chickadees & two Kinglets more in evidence now.

Melospiza melodia morphna

Thryomanes bewicki calophonus

Spinus tristis salicamans

Colaptes cafer cafer

Spinus pinus pinus

Turdus propinquus saurinus. Flock tendency one 30 birds

Pipilo maculatus oregonus

Buteo borealis calurus.

Junco oregonus shefeldti.

Hepocichla ustulatus ustulatus.



It is every evident that many of the more familiar birds of summer are gone and a slight fall change in other forms.

Took pictures (1-2-3 10-4-42) of environs of Severeth Creek and adjoining bench.