

*Anas albifrons albifrons*

*Branta canadensis*. (slightly smaller than white-fronts)

*Squatarola squatarola*.

*Falco peregrinus pealii*. Extremely black, but found that the Canadian and white-fronts could not be told apart with plumage design being entire black under the present lighting conditions. Likewise this hawk may have appeared darker than usual.

*Gus canadensis tabida*

*Fulica americana*

*Nettion carolinense*.

*Mareca americana*

*Chaulelasmus streperus*.

*Ardea herodias fannini*

*Passerculus sandwichensis alaudinus*

*Euphagus cyanocephalus*

*Melospiza melodia morphna*

*Larus occidentalis*.

*Oxyechus vociferans vociferans*

*Colaptes cafer cafer*.

*Sturnella neglecta*

*Phasianus colchicus torquatus*

*Spinus tristis salicamans*

*Myroca marila*.

*Phalacrocorax auritus albicollatus*.

*Sayula acuta tzytzehaa*.

*Anthus sprusletta rubescens*.

*Mergus merganser* (Sp.)

At sundown as I was leaving the western Vancouver pasture area observed 82 white-fronted trending south in same direction as the Sandhills had previously taken. This group was flying high and in earnest as to destination.

[see page 420509-40 for photo dated Oct 11, 1942]

noticeable signs and feeling of autumn. Wind from the east which is a good sign and generally indicative of a change of season.