

5/15/43.

Census of Lewisish and Vancouver lake area. Itinerary as of trip of 5/8/43. In field from 1:00 P.M. to advanced twilight or approx 8:00 P.M. weather predominately rain with intermittent periods of sunshine and blue patches of sky. Day cold with respiration discernible in afternoon. Forms observed.

Sturnella neglecta - approx. 13

Passer domesticus - residential.

Spizella passerina arizonae - 6

Stelgidopteryx thalassina lepida - Generally represented in entire area traversed. One group persistently controlling an area of creek bottom continued to patrol same area in slightly diminished numbers throughout one of the short showers. Their greatest activity preceded the shower proper.

Icteria virens longicauda. 5 isolated individuals. calling continually. One observed to leave its protective covering and invade the grasses at peripheral edge of the bush patch.

Dendroica aestiva brewsteri approx. 18. Observed certain degree of coaction between this form and the Helminthophila pusilla. Called occasionally.

Turdus migratorius caurinus. Generally represented

Phasianus colchicus torquatus.

Melospiza melodia morphna. These forms are never as conspicuous or in evidence as during the winter. Their unquestonable due to protection of foliage coverage. They were heard frequently in song.

Zonotrichia leucophrys pycnonis. Heard throughout day and observed very few otherwise.

Carpodacus mexicanus frontalis. Heard very frequently throughout the day. Cassini more commonly represented however.

Spinus tristis salicemans. Generally represented. Heard one sing with a respectable volume and quality.

Helminthophila pusilla. Approx 12.

Geothlypis trichas occidentalis. Approx 7. Have never found them associated with water courses but among the underbrush of sidehills where they are associated with the cat, towhees and warblers.