

Columbo fuscata fuscata. 12 together on top of fir tree in area of semi- virgin timber. This was shortly after the shower and their positions indicated that they were utilizing the sun rays as they were found on the very top of the tree. They left <sup>at 30'</sup> and flew across the canyon 500' away and lit on top of another fir.

Chaetura vociferi. Approx 20. in small groups.

Psaltiriparus minimus minimus. Heard one bird

Cyanocitta stelleri stelleri. Group 2 birds.

Air sparrow. Only common native bird represented by 24 birds in small groups. 6 mallards only other birds observed.

Ardea herodias farrini Approx. 12. most of these birds very large.

Passerculus sandwichensis. Commonly represented in field with a wild guess of about 250 birds.

Actitis macularia. 4 birds of 2 pair.

Larus glaucescens.

Larus delawarensis. (?) Have never observed so few gulls in evidence as today. Only 2 group of 8 birds flew over lake in their evening movement from Columbia river. 12 other gulls observed.

Otocoris alpestris. Observed three but am sure that there were many represented in the fields with possibly the Pipit mixed in among them.

Agelaius leucis. 1 bird only in same area as of last week.

Pluvialis dominica. Satisfied myself twice on the identification of this bird and under most starting and favorable conditions. Only questionable fact being that the black extended only to region of legs and not to base of tail. Tail and upper coverts brown and marked like back, no <sup>black</sup> auxiliary feathers, dark head, wing same intensity of coloring. This bird reacted different than the black bellied plover. When approached during the initial stalk, it would respond by elevating entire body, neck and head as a rigid unit, tilting slightly backward. The movements were not as dignified