

weather condition for bird census than an even slightly windy day. When leaves are moving it is difficult to pick up the movements of birds among trees and shrubs. In field from 1:00 P.M. to 8:30 P.M. Recorded the following forms: (all counts approximate.)

Passer domesticus. residential

Turdus migratorius caurina. usual

Piranga ludoviciana. 6

Stelgidopteryx thalassina lepida usual

Iredoprocne bicolor. usual

Hedymeles melanocephalus papago. 4

Hylaechthya ustulata 4

Dendroica aestiva. 20

Icteria virens longicauda 6

Penthestes atricapillus oregonus 4

Psaltriparus minimus minimus 12 One family of 5 following adults.

Vireo solitarius cassinii. 7 One in dense conifer, area with call like wren and deciduous

Spyzella passerina arizonae. 10

Melospiza melodia morphna. usual

Oryzopsis vociferans vociferans. 28. Evidence of forming aggregations of 8-10.

Pipilo maculatus oregonus 10

Passerina amoena 3

Hirundo erythrogastrer. Usual representing hundreds and more or less found in greater number along lake shore in contrast to the occurrence of N.V. grays and tree swallows found more inland and associated with conifers etc.

Sporus tratis sullivanensis. 18. Small groups.

Zonotrichia leucophrys purgetensis. 12

Geothlypis trichas occidentalis 8

Calaptes cafer cafer. 3. noticeable decline in numbers

Bombicilla cedrorum. 25. Small groups - Have been no evidence for last week.

Dendroica auduboni 4

Vireo gilvus swainsoni 8