

Nyctanassa guttata - 2Phasianus colchicus torquatus - usualDendroica nigrescens 2. In same area as observed on previous trips.Oporornis tolmiei 1Regulus satrapa olivaceus 2Columba fuscata fuscata 1Selasphorus rufus. no record in upper Liverich but in group of 6 with aggregate of 15 associated with new flowering slopes of railway grade near lower limits of the creek. They were definitely associated with their suitable habitat.Vermivora celata. 1 Characteristic call. +kill → ! ! ! ! ! ! ! !Ardea herodias farrumii 14 in south end of lake.

Evidence of territorial desires of each bird. Some larger birds aggressive and with out others along shore line. Birds feeding in recently inundated areas.

Aix sponsa 8 fewer in evidence.Corvus brachyrhynchos hesperis. 6 - Decided depreciation of numbers. They might shift in position according to high water levels as they were not in this area today. An area normally supporting great numbers of these birds!Icterus bullocki 1. mouth LiverichJunco oregonus oregonus 4 - depreciation of numbers.Carpodacus mexicanus frontalis. 15Zenaidura macroura macroura. 2Otocoris alpestris. 9Euphagus cyanocephalus. 3Anas platyrhynchos platyrhynchos 4Tyrannus vociferans 2Passerculus sandwichensis alaudinus 12Sturnella neglecta - 3Buteo borealis calurus 1Sitta carolinense. 1Larus glaucescens 5Chordeiles minor 1

(12 unknowns passed on at twilight in usual flight lane)

In general the bird life was affected by the change of water level of the lake. Expected to find an aggregation of birds around