

Colaptes cafer cafer - several groups of ^{3 + 4}
Corvus brachyrhynchos brachyrhynchos. Collecting in large groups. Direction of casual crow flight generally indicates direction of congregation. Vesperous calls can be heard from some distance.

Melospiza melodia morphna - several singing and defended territories

Corthylio salindula gunnelli

Zonotrichia coronata. 3.

Anas platyrhynchos platyrhynchos

Chaulelasmus streperus 2

Mareca americana. 600. This species frequently found resting on land surface in large flocks some 50 to 200 feet from water. When approached generally leave with single impulse. Dominant species

Oxyechus vociferans vociferans - 18

Nettion carolinense. 500. Call single as contrasted to double call of Baldpate.

Buteo borealis calurus. 1

Regulus satropa olivaceus

Capela acuta tzitziloo. 130

Pipilo maculatus oregonus 3

Thryomanes bewicki calophonus 2

Tulica americana 12

Dryobates pubescens gardneri 2

Ardea herodias farreri 15

Branta canadensis 4

Spatula clypeata 20

Nyroca valisineria 8

Horned lark or pipit 1

Haliaeetus leucocephalus alaskanus 1 immature.

Phalacrocorax auritus 8

Larus glaucescens

Larus brachyrhynchos. Counted all gulls of the various species intercepting line across south end of Vancouver lake on the northern afternoon flight. 176 gulls in 5 minutes. Time 6:30 PM at about 20 minutes before sundown.

Phasianus torquatus calchasi 4

Nycticorax nycticorax - ? 1

Euphagus cyanocephalus

Aechmophorus occidentalis 1

Passerella iliaca townsendi 2