

Ardea herodias tringoides. One bird at Mono Lake.

Anas platyrhynchos platyrhynchos. Few at Mono Res. also represented at Chukew and Lower Bridge Res.

Arca piscivora hudsonius. Only 4 birds observed.

Buteo borealis calurus: Specimen procured across Lewon bench from highway. Measured wing $15\frac{1}{2}$ " , tail 8 inches, tarsus $3\frac{3}{4}$ inches, beak $1\frac{3}{8}$ inches, length $20\frac{1}{2}$ inches, wing spread $46\frac{1}{2}$ inches. Other birds observed circling in air.

Citellus grammurus. First one observed at Lewon set. on top of fence post. Others occasionally observed to south.

Sialia currucoides. From Nephis south to Kanash the bluebirds increased in numbers until they reached their maximum in numbers at the Eagle nest area. An ordinary count would be approx 8 birds per mile. Male and females represented. Generally found sitting on fence line and telephone lines.

Corvus corax sinuatus. The bird of the day. This bird was found generally distributed from Nephis south to Kanash increasing in percent frequency toward the south. Recorded groupings as 1-3-2-3-3-10-3-2-4-3-3-3-6-1-1. On return from Kanash found one group of 37 birds which were feeding at a range cattle feeding pen at north edge of Kanash. These birds were generally found feeding in wheat field or soaring in air. Two rooves in quartzite outcroppings south and west of Kanash. No nesting activity observed.

Circus hudsonius 5 birds (singles) at reservoirs.

Nyroca valisineria. Few birds at Chukew Creek Res.

Fulica americana few birds at Chukew Creek Res.

Nettion carolinense Few birds at Chukew Creek Res.

Nyroca americana. Approx 200 feeding in Chukew Creek Res.

Querquedula cyanopectus. Approx. 35 in one group at Chukew Creek Reservoir. At the lake there were possible 2000 ducks on this reservoir, most of them probably redheads.

Mergus mercator americana. (Sp ?) could be redbreasted. 18 in Lower Bridge Reservoir. They were feeding in one group when