

Other birds shot near this trapline are:

- 11-3-26-48 *Melospiza melodia*  
 12-3-26-48 Mockingbird  
 13-3-26-48 White-throated sparrow  
 14-3-26-48 " " "  
 15-3-26-48 Brown Creeper  
 16-3-26-48 Audubon Warbler  
 17-3-26-48 *Melospiza melodia*  
 18-3-26-48 White-throated sparrow  
 19-3-26-48 " " "  
 20-3-26-48 " " "

noted also one copperhead snake, 1 brown thrasher, black-capped chickadee, towhee, blue jay & turkey vultures.

The 100 traps set in damp pasture of sedge & grass did not yield a single animal although I judge this field to be the most productive. Rain damage may have been the main factor as a greater percentage of traps were sprung.

From the 83 traps set along forest edge produced the following:

- 21-3-26-48 *Reithrodontomys* 160-90-20-14-10gms testes 8mm.  
 22-3-26-48 " 170-95-21-12gms Testis 8mm  
 23-3-26-48 *Pitymys nemoralis* 129-26-18-11.5-30gms no embryos  
 24-3-26-48 " " 89-19-15-8-8gms ♂  
 25-3-26-48 " " 135-28-18-11.5-46gms (2 x 1 emb. 26mm)  
 26-3-26-48 " " 135-27-18.5-11-31gms ♂ testes 8mm  
 27-3-26-48 " " 105-24-17-10-15gms ♂  
 28-3-26-48 " " 102-26-16.8-10.5-15gms ♀

### Mena, Arkansas

March 27, 1948

Last night extremely cold with high march wind. morning cold  
 Inspected line of 60 traps and caught the following:

- 1-3-27-48 *Pitymys nemoralis* 133-29-19- ♂ skull only, testes 9mm  
 2-3-27-48 Swamp sparrow. caught in trap.  
 3-3-27-48 *Pitymys nemoralis* 118-26-17. ♂ skull only, testes 5mm  
 4-3-27-48 *Sturnella neglecta* caught in trap.

From set of 83 along edge of forest & open grass field caught:

- 5-3-27-48 *Pitymys nemoralis* 118-24-16.5-11-16gms ♂  
 6-3-27-48 " " 120-25-16.5-10-23gms ♂ testes 5 1/2 mm  
 7-3-27-48 *Reithrodontomys* 138-75-19-14-8gms (no embryos)  
 8-3-27-48 " 145-77-19-14-8gms testes 4mm