

*Selasperus platycircus*, *Scondroica auduboni*, *Nuttallornis borealis*, *Carpodacus cassinii*, *Passercetes grammurus*, *Loxia curvirostris*, *Buteo jamaicensis borealis*, *Accipiter gentilis*.  
Returned to camp at 9 mi. S Robertson and finished preparation of mammals.

9 mi. S Robertson, 8400 ft., Uinta Co., Wyoming.

July 5, 1948

Inspected trapline of research area A-7-4-48 of last evening setting:

1. sprung; 2-4 uneffected; 5 sprung; 6-9 uneffected; 10-11 sprung; 12-14 uneffected; 15 sprung; 16 *Peromyscus maniculatus* 1-7-5-48; 17-18 uneffected; 19 *Bufo* 2-7-5-48; 20-25 uneffected; 26 *Microtus longicaudus* 3-7-5-48; 27 uneffected; 28 *Zapus princeps* 4-7-5-48; 29-32 uneffected; 33 sprung; 34 *Microtus longicaudus* 5-7-5-48; 35-36 uneffected; 37 *Eutamias umbrinus* 6-7-5-48 from mud flat of receding edge of lake; 38 sprung; 39-40 uneffected; 41 *Zapus princeps* (tail only); 42 sprung; 43 *Eutamias umbrinus* 7-7-5-48 on mud of receding lake shore; 44-53 uneffected; 55-58 uneffected; 59 sprung; 60-65 uneffected; 66-67 sprung; 68 *Melospiza lincolni* 8-7-5-48; 69-70 uneffected; 71 *Microtus longicaudus* 9-7-5-48; 72-73 sprung; 74 uneffected; 75 sprung; 76 *Melospiza lincolni* 10-7-5-48; 77 *Zapus princeps* 11-7-5-48; 78 uneffected; 79 *Microtus montanus* 12-7-5-48; 80 *Microtus montanus* 13-7-5-48; 81-83 uneffected; 84 sprung; 85-89 uneffected; 90 sprung; 91-94 uneffected; 95 sprung; 96-110 uneffected; 111 sprung; 112-113 uneffected; 114 sprung; 115-117 uneffected; 118 sprung; 119 sprung; 120-120 uneffected; 121 sprung; 122 uneffected; 123 *Peromyscus maniculatus* 14-7-5-48; 124 *Eutamias umbrinus* 15-7-5-48; 123-125 uneffected; 126 *Peromyscus maniculatus* 16-7-5-48; 127-137 uneffected; 138 sprung; 139 uneffected; 140 sprung; 141 uneffected; 142 *Peromyscus maniculatus* 17-7-5-48; 134 uneffected. The above series from 83-143 eaten by ants to some degree; 144 uneffected; 145 uneffected; 146 *Zapus princeps* 18-7-5-48; 147 uneffected; 148 *Microtus montanus* 19-7-5-48 taken 2 feet from 18-7-5-48; 149-153 uneffected. In summary: *Microtus montanus* and *M. longicaudus* will enter drier areas that have overhead protection of willows & grasses but will not enter coniferous or aspen forests even where they are adjacent to lake grasses. Returned to camp and then to Gilbert meadows, Summit Co., Utah, 9641 feet some 7 miles up canyon from camp. Enroute passed thru extensive logging operations about 1/2 way. Considerable trees