

supplied with water from spring some 150 feet inland. Photo 22-7-14-48 in same area with rocks in foreground where beaver fed in evening upon willows. Elevation at this point approx. 10,500 ft. Returned to Brush Creek Camp and by 11:30 A.M. had organized for departure to Hayden Range to the west. The itinerary will include Riverside Encampment, thence west on forest service road to point 8 mi N and 19 1/2 mi. E Savery, Carbon Co., Wyoming at Lost Creek Camp Grounds at 8800 ft. At 5 mi. N on highway 230 from Riverside observed 2 *Aquila chrysaetos canadensis* along the east side of road at a point where it passes near the rocky outcroppings. At the moment I saw them they were in the air above the ledges. The lower bird was an immature and carrying a *Citellus richardsoni*. The adult was 20' above it and watched its every movement. The young called continuously. They soared and circled several minutes and then alighted on one of the rocky outcroppings. The young continued to play with the *Citellus*. The adult, 10 feet away watched it. After 5 minutes they both left and continued on to the south, the young still carrying the *Citellus*. This behavior is in contrast to the retiring nature of eagles during nesting time - their main concern at that time is to get out of the country without being seen. From the topography of the area would judge that the birds had used a nest in the immediate area. Continued south thru Riverside and Encampment, thence west to divide of Hayden Range. At 8250 ft and approx. 5 mi. west of Savery, observed a *Lepus townsendi* cross the road and take its position by a culvert. When car was stopped, it remained until 10 feet at which time it left, ran 30 feet and stopped again. Continued west to divide where I noticed considerable lumbering operation. At this point there is considerable open areas. A rock slide on north ridge should produce *Ochotona*. There is something about Hayden Range that is different from other ranges of equivalent height (our elevation) and plant matrix. It may have suffered overgrazing or extensive timber cutting or fire or possibly all three. Continued down west drainage and camped at Lost Creek Camp Grounds at 8 mi N + 19 1/2 mi E Savery, 8800 ft., Carbon Co., Wyoming. Set series of traps near camp. Research area A-7-14-48 among heavy grasses and vegetation bordering creek, open but bordered by willow, aspen and spruce. This community should be favorable for *Zapus*