

dominant grasses: 30-8-27-48; 31-8-27-48; 31(1)-8-27-48;  
31(2)-8-27-48; 31(3)-8-27-48; 31(4)-8-27-48; 31(5)-8-27-48. Continued  
W to Blackfoot, Montana Set 120 traps.

4 mi. E and 10 mi. S Blackfoot, 3900ft., Glacier Co., Montana  
Aug 28, 1948

Caught 15 mammals in 120 traps. Grasses 16(1)-8-28-48;  
16(2)-8-28-48; 17(1)-8-28-48; 17(2)-8-28-48; 17(3)-8-28-48; 17(4)-8-28-48;  
17(5)-8-28-48; 17(6)-8-28-48; 17(7)-8-28-48; 17(8)-8-28-48; 18-8-28-48;  
19-8-28-48. Continued to Springdale, set 120 traps.

Springdale, 4100 ft., Park Co., <sup>Montana</sup> Wyoming  
Aug 29, 1948

In 120 traps caught 60 mammals (see catalogue). Dominant  
grass 70(1)-8-29-48 and 70(2)-8-29-48. Field 80% cut & prone.  
nos <sup>grasses</sup> 71(1)-8-29-48; 71(2)-8-29-48; 71(3)-8-29-48; 71(4)-8-29-48;  
71(5)-8-29-48 and 71(6)-8-29-48 used by meerkats. Continued  
to Greybull and set traps in several places.

8 1/10 mi. S Greybull, 3788 ft., Bighorn Co., Wyoming  
Aug 30, 1948

\* From 10 traps took two *Microtus ochrogaster* nos 1-8-30-48  
and 2-8-30-48. Dominant grasses 6(1)-8-30-48; 6(2)-8-30-48;  
7(1)-8-30-48; 7(2)-8-30-48.

1 mi. S Greybull, 3795 ft., Bighorn Co., Wyoming  
Aug 31, 1948

From 10 traps took 7 *Mus musculus*. no 8-8-30-48 dominant  
grass.

7 1/2 mi E Greybull, Bighorn Co., Wyoming  
Aug 31, 1948

From 10 traps collected 1 *Reithrodontomys* 4-8-30-48.  
Several traps were set in good runways of *Microtus ochrogaster*  
but all snopped & without mammals.

\* [31 mi. E and 10 mi. <sup>(?)</sup> W Greybull, Bighorn Co., Wyoming  
Aug 30, 1948

Collected one *Tamiasciurus*.]

From Greybull continued S. *Sylvilagus* 1 per 10 miles  
of road kill. Golden eagle 20 mi. S Buffalo. Area  
near Midwest appears to be formidable to *Microtus ochrogaster*.