

1 1/2 mi. W and 2 1/4 mi. S Casper, 5250 ft., Natrona Co., Wyoming. 8 Aug, 1948.
Grassland-Transition contact. Entrenched creek at base of range.
Grasses from erosional level one foot above the present creek level. Soils slightly damp.

5(1)-8-6-48 (dominant)
5(2)-8-6-48 "
6-8-6-48 "

7 mi. S and 2 mi. W Casper, 6300 ft., Natrona Co., Wyoming. 8 August, 1948.
Grassland-Transition contact. Isolated willow stand with spring water forming a damp meadow below. On piedmont of grassland bordering range.
Aspen and Ponderosa Pine associated.

11-8-8-48 (dominant)
12-8-8-48 (dominant)

6 mi. W. and 2 mi. W Casper, 5900 ft., Natrona Co., Wyoming. 9 August, 1948.
Grassland-Transition contact. Wet meadow below spring area on piedmont.

5(1)-8-9-48 (dominant)
5(2)-8-9-48 "
5(3)-8-9-48 "
6(1)-8-9-48 "
6(2)-8-9-48 "

3 2/10 mi. E and 6/10 mi. S Cody, 5020 ft., Park Co., Wyoming. 12 August, 1948.
Spring and marsh on benchland in creek valley.

40-8-12-48 (dominant)
41(1)-8-12-48 "
41(2)-8-12-48 "
42(1)-8-12-48 (subdominant)
42(2)-8-12-48 "
42(3)-8-12-48 "

1 mi. W and 8/10 mi. S Buffalo, 4600 ft., Johnson Co., Wyoming. 17 August, 1948.
Grassland-Transition contact. Marsh in river valley. Considerable alkali and mud holes associated.

13-8-17-48 (dominant)
14(1)-8-17-48 "
14(2)-8-17-48 "
14(3)-8-17-48 "
15(1)-8-17-48 (subdominant)
15(2)-8-17-48 "
15(3)-8-17-48 "
15(4)-8-17-48 "
15(5)-8-17-48 "
15(6)-8-17-48 "
15(7)-8-17-48 "
16(1)-8-17-48 (dominant)
16(2)-8-17-48 "

6 1/2 mi. W and 1 mi. S Buffalo, 5600 ft., Johnson Co., Wyoming. 18 Aug., 1948.
Grassland-Transition contact. Canyon floor at edge of creek. Grasses completely covering area. Wild rose, gooseberry, chokecherry associated nearby. Hillside of Artemisia and Pinus ponderosa and dry grasses.
Canyon crowded with cottonwoods, willow and typical stream side vegetation.