

The species of mammals collected are all considered to be common to the general area and include the following:

Microtus ochrogaster ochrogaster (Wagner)
Sigmodon hispidus texianus (Audubon and Bachman)
Peromyscus leucopus noveboracensis (Fisher)
Peromyscus maniculatus bairdii (Hoy and Kennicott)
Reithrodontomys sp?
Mus musculus Linnaeus.
Pitymys nemoralis (Bailey)

Frequency and percent frequency of the various members of the species in the aggregate of the research areas.

<u><i>Microtus</i></u>	<u><i>Sigmodon</i></u>	<u><i>Peromyscus L</i></u>	<u><i>Peromyscus m.</i></u>	<u><i>Reithro</i></u>	<u><i>Mus</i></u>	<u><i>Pitymys</i></u>
66 - (25%)	126 - (47%)	7 - (.03%)	47 - (17%)	18 - (07%)	1 - (001%)	1 - (004%)

Frequency and percent frequency of species in individual research areas.

	<u><i>Microtus</i></u>	<u><i>Sigmodon</i></u>	<u><i>Peromyscus L</i></u>	<u><i>Peromyscus m.</i></u>	<u><i>Reithro</i></u>	<u><i>Mus</i></u>	<u><i>Pitymys</i></u>
A	21 - (24%)	46 - (52%)	7 - (.08%)	9 - (17%)	5 - (05%)	0 - (0%)	0 - (0%)
B	21 - (22%)	46 - (50%)	0 - (0%)	19 - (20%)	8 - (08%)	0 - (0%)	0 - (0%)
C	4 - (15%)	10 - (37%)	0 - (0%)	11 - (40%)	1 - (03%)	1 - (03%)	0 - (0%)
D	20 - (35%)	20 - (42%)	0 - (0%)	8 - (12%)	4 - .14%	0 (0%)	1 - (02%)

Visitation to individual trap sets during 7 day period.

Area A	Area B	Area C	Area D
197 (56%)	214 (61%)	49 (14%)	143 (41%)

Individual trap visitations.

Area A	Area B	Area C	Area D
47 - (94%)	48 (96%)	30 (60%)	47 (94%)

Visitation of individual traps as an aggregate. 172 - (86%)

Sex ratio of the species in the aggregate.

<u><i>Microtus</i></u>	<u><i>Sigmodon</i></u>	<u><i>Peromyscus L</i></u>	<u><i>Peromyscus m.</i></u>	<u><i>Reithro</i></u>	<u><i>Mus</i></u>	<u><i>Pitymys</i></u>
♂ 27 (41%)	57 (45%)	5 (71%)	25 (53%)	7 (39%)	0 (0%)	0 (0%)
♀ 39 (59%)	69 (55%)	2 (29%)	22 (47%)	11 (61%)	1 (100%)	1 (100%)

Sex ratio in individual areas.

	<u><i>Microtus</i></u>	<u><i>Sigmodon</i></u>	<u><i>Perom. L</i></u>	<u><i>Perom.</i></u>	<u><i>Reithrodontomys</i></u>	<u><i>Mus</i></u>	<u><i>Pitymys</i></u>
A	♂ 8 (38%)	21 (45%)	5 (100%)	5 (55%)	1 (25%)	0 (0%)	0 (0%)
	♀ 13 (62%)	25 (54%)	0 (0%)	4 (45%)	3 (75%)	0 (0%)	0 (0%)
B	♂ 8 (38%)	19 (41%)	0 (0%)	11 (58%)	3 (37%)	0 (0%)	0 (0%)
	♀ 13 (62%)	27 (59%)	0 (0%)	8 (42%)	5 (63%)	0 (0%)	0 (0%)
C	♂ 2 (50%)	6 (60%)	0 (0%)	5 (55%)	2 (100%)	0 (0%)	0 (0%)
	♀ 2 (50%)	4 (40%)	2 (100%)	6 (45%)	0 (0%)	1 (100%)	0 (0%)
D	♂ 9 (45%)	11 (46%)	0 (0%)	4 (50%)	1 (25%)	0 (0%)	0 (0%)
	♀ 11 (55%)	13 (54%)	0 (0%)	4 (50%)	3 (75%)	0 (0%)	1 (100%)