

## JOURNAL 1949

Museum of Natural History, University of Kansas, Lawrence, Kansas

Jan 2, 1949

This morning skinned out some Coyote-dog hybrids. Recorded several black and white photographs of these hybrids. Photos 1-1-2-49 to 12-1-2-49 as follows:

- (1-1-2-49) ♂ *Canis latrans* (hybrid) full view. Stomach empty. Measured 1340-390-216-47 lbs. (see Hall no. 6760)
- (2-1-2-49) ♂ *Canis latrans* (hybrid) full view. Cow hair in stomach. (see Hall field no 6761)
- (3-1-2-49) ♂ *Canis latrans* (hybrid) full view. Pig hair in stomach. Also feathers flakes of skin as if from butchering of a hog (see Hall no. 6762)
- (4-1-2-49) ♂ *Canis latrans* (hybrid). Head view only (see Hall no 6761)
- (5-1-2-49) ♂ *Canis latrans* (hybrid) Tail only showing bushy condition (see Hall no 6761)
- (6-1-2-49) ♂ *Canis latrans* (hybrid) Hind foot only (see Hall no 6760)
- (7-1-2-49) ♀ *Canis latrans* (typical) Full view (see " " 6763)
- (8-1-2-49) ♀ *Canis latrans* (" ) head view only (see Hall 6763)
- (10-1-2-49) ♀ *Canis latrans* (" ) hind foot only (see Hall 6763)

These Coyotes were collected on Dec 31, 1948 by Edward A. Revey at 2 miles south of Le Laupé, Franklin Co., Kansas. The typical Coyotes were also collected by Revey on Jan 2 (this date) 1949 in the same general area as was the case of the capture of the hybrid Coyotes. Photo 11-1-2-49 in color of *Canis latrans* (hybrid). Same as photo 2-1-2-49 of above.

The following photographs recorded in notes are on the following page:

- 4-1-2-49 of hybrid showing similarity to dog in profile
- 8-1-2-49 a typical Coyote profile, hair darker.
- 2-1-2-49 hybrid showing shorter legs & heavier flanks in hind quarters.
- 7-1-2-49 typical Coyote
- 6-1-2-49 hybrid foot - relatively wide like in a dog
- 10-1-2-49 typical Coyote with narrow foot