


Climacodum. Small <sup>large</sup> form only. 490721-35  
Solaster stimpsoni 1 specimen rare dead shell of larger  
Corpidula. Small Corneal forms  only.

Stichopus californicus 3 specimens  
Modiolus modiolus few unbedded ~~specimens~~ securely  
in matrix of debris of algae halberds and other material.  
appeared to be encased in a slimy covering.

4 or 5 spec. of shrimp Few white worms.  
Pagurus heunerlyi. few. 2 spec. of Isopods - numerous  
Neptuna tuberculata. dead shells only

Pugettia gracilis & most common form but not too  
well represented. Form from gray to red in color.

Cucumaria. Small white form.

Small sculpin. 3.

Three species of blenny, one of which was green.

Epiactis productus 1

1 nudibranch small and filamentous decoration.

Cryptostomus. 27 cm in length.

Ischnochelone. small in size and 5 collected. and others.

Nereocystis, Laminaria, Desmarestia, Costaria represented  
and constituted 80 percent of bulk

Lacuna common on help.

Coprella few. 1 Duck shell (shell only) measuring

This dredge as stated above supported several forms of  
coral. as a result of position away  
from immediate outlet of False Bay channels.

Station 6-7-21-49 differed from above in a noticeable lack  
of the coral representatives and a slightly greater number  
of S. frutescens. The period of dredge was about  $\frac{1}{6}$  of  
the dredge 5-7-1-49. Algae occurred in about the same  
percentage of hauls.

Station 2-7-21-49 supplied the greatest number of S. frutescens  
10 in number and constituted  $\frac{1}{4}$  of the bulk. 3 Stichopus  
californicus represented. This station approached station  
5-7-21-49 in position but did not have the coral representation  
to the same degree. In addition to the above, <sup>listings off other dredges</sup> took a  
small brownish fish probably Ascelichthys rhodurus. Also 2 Cancer  
oregonensis. One large Polonicea. One Euphrosia. 35 m.m long

Enroute to False Bay recorded the following birds. These  
birds were all observed near Cattle point and channel to  
east around Hall islands.

Puffinus griseus. Two birds left water at 400 meters  
east and south of cattle point.