

| | | | | |
|------|------------|---|------------------------------|------------------------------------|
| m.o. | 510817-99 | ♂ | <i>Clethrionomys rutilus</i> | 125-34-19-13-20 gms, ut. normal |
| " | 510817-102 | ♀ | " | 131-34-19-13-19 " , " " |
| " | 510817-103 | ♂ | " | 130-33-18-13-20 " , testes 2.5 mm |
| " | 510817-104 | ♀ | " | 130-33-19-14-20 " , " " |
| " | 510817-105 | ♂ | " | 123-36-19-14-20 " , testes 3 mm |
| m.o. | 510817-106 | ♂ | " | 130-34-19-14-21 " , " 2.5 " |
| " | 510817-107 | ♀ | " | 128-33-19-13-18 " , ut. normal |
| " | 510817-108 | ♂ | " | 126-33-19-13-28 " , testes 2.5 mm |
| " | 510817-109 | ♂ | <i>Microtus murus muriei</i> | 110-22-17-12-13 " , |
| " | 510817-110 | ♂ | <i>Spizella arborea</i> | L. 160, 18 gms |
| " | 510817-111 | ♀ | <i>Microtus oeconomus</i> | 180-48-20-13-65 gms, 2x4 emb 21 mm |
| " | 510817-112 | ♀ | " <i>murus muriei</i> | 133-29-19.5-13-27 " , ut. normal |
| m.o. | 510817-113 | ♀ | <i>Spizella arborea</i> | L. 156, 18 gms |

Aug. 18, 1951

| | | | | |
|-------|------------|---|--|--|
| | 510818-1 | ♂ | <i>Spermophilus u. kennicottii</i> | 385-122-63-13-620 gms, testes 9 mm |
| | 510818-5 | ♀ | <i>Clethrionomys rutilus</i> | 152-41-19-14-33 gms 6x6 plac scars |
| | 510818-15 | ♀ | " | 143-43-20-14-38 " , lactating |
| | 510818-34 | ♂ | <i>Microtus murus muriei</i> | 121-27-20.5-13-24 gms, testes 3 mm |
| m.o. | 510818-49 | ♀ | <i>Clethrionomys rutilus</i> | 112-34-18-13-18 gms, ut. normal |
| " | 510818-51 | ♂ | <i>Microtus murus muriei</i> | 120-28-20-13-19 " , testes 2.5 mm |
| | 510818-55 | ♂ | " | 128-28-21-13-24 " , " 3 " |
| | 510818-59 | ♂ | " | 131-34-21-13-23 " , testes 3 mm |
| m.o. | 510818-73 | | <i>Clethrionomys rutilus</i> | 118-33-18-13-15 gms. |
| | 510818-77 | ♂ | " | 124-36-19-13-17 gms, testes 2.5 mm |
| m.o. | 510818-78 | ♀ | <i>Microtus murus muriei</i> | 100-24-18-12-11 gms, ut. normal |
| | 510818-88 | ♀ | " | 88-33-20-14-39 " , 4x2 plac. scars |
| | 510818-91 | | <i>Clethrionomys rutilus</i> | 126-36-20-15-22 " , |
| m.o. | 510818-92 | ♂ | <i>Spizella arborea</i> | L. 156, wt 20 gms |
| " | 510818-100 | | <i>Clethrionomys rutilus</i> | 140-43-20-14-16 gms |
| | 510818-101 | ♂ | <i>Microtus oeconomus</i> | 170-44-20-13-50 " , testes 5 mm |
| m.o. | 510818-104 | ♂ | " | 105-28-17-11-12 gms, testes 4 mm |
| | 510818-112 | ♀ | <i>Sorex cinereus agyrak</i> | 98-34-12-7-3.5 gms, ut. normal |
| m.o. | 510818-115 | ♂ | <i>Microtus oeconomus</i> | 115-32-19-12-15 gms, |
| | 510818-121 | ♂ | <i>Clethrionomys rutilus</i> | 118-31-19-14-21 gms, testes 2.5 mm |
| | 510818-124 | ♂ | <i>Microtus oeconomus</i> | 168-45-21-13-52 gms, testes 7 mm |
| | 510818-125 | ♂ | " | 152-42-20-12-42 " , " 6 " |
| | 510818-127 | ♀ | " | 146-38-19-13-42 " , plac. scars & vascular |
| m.o. | 510818-128 | ♀ | <i>Clethrionomys rutilus</i> | 127-33-19-13-20 gms, ut. normal |
| | 510818-129 | ♂ | " | 120-30-19.5-13-24 " , testes 3 mm |
| | 510818-150 | ♂ | " | 120-30-19.5-13-21 " , testes 2.5 mm |
| skel. | 510818-151 | | <i>Calcarurus l. alascensis</i> (juv). | L. 170 mm, wt 23 gms. |