



with shallow excavations. The mice had reestablished runways and had dug new runways thru the freshly excavated dirt. Observed one *Microtus micurus* active at 11:20 A.M., day clear. They have the life form of the pikas. Wind down Canyon for the first time. Specimen 520802-11 *Lemmus* and 520802-12 *Microtus micurus* from grass-sedge refugia and taken shortly after setting. Today at 11:30 P.M. these same traps were not effected by other mice. From 4:00 P.M. to 5:00 P.M. wind gradually subsiding. At 5:30 P.M. to 7:00 P.M. wind increased to this A.M. as an intense wind. Golden eagle eating dead caribou this morning on delta or 2nd alluvium down east shore from S end Lake Peters. Observed an Am golden eagle also up canyon. These eagles favor the Lake <sup>Peters</sup> ~~Schneider~~ area and are frequently are seen at all times of the day probably hunting marmots. This predation might account for the weerness of the marmots. One large dark bird and one smaller one with this degree of white in tail  flew and sailed in high wind on the NE face of mount Mary at 8:30 P.M. thus:

The method of hunting was to circle up steep canyons to head of canyon and then drop down next canyon to surprise animals at bottom of canyon and the entrance of the next adjoining canyon. The same act is repeated on down the slope of the mountain range. Temp. at midnight 52°F, min 32°F, max. 70°F.



Aug. 3, 1952

From trapline collected the following:

520803-1 *Lemmus* from refugia area of yesterday trapping indicating some mice are still wandering around after having been displaced by caribou trampling.