

Troops placed ^{10' apart} mainly in damp to wet meadows of grasses and sedges and willow. Some at base of rocks at edge of the meadow and a few on higher ground among aspens on damp soils. The arm of the lake is filling in and the outer region is tall sedges & grasses; inland it is shorter grasses and sedges with willow mixed in. From traps collected the following:

- 21 adult and 64 subadult of *Microtus pennsylvanicus*
- 4 *Synaptomys* (1 of which is a subadult).
- 1 *Sorex palustris*
- 2 *Peromyscus*
- 2 *Eutamias*
- 16 *Zapus*
- 13 *Clethrionomys gapperi*
- 4 Lincoln sparrows
- 1 golden-crowned sparrow
- 4 song sparrows
- 1 garter snake.
- 8 *Rana pipiens*

Microtus pennsylvanicus from marsh sets. Two of the *Synaptomys* from 10 feet apart.

A few generalizations are:

Microtus pennsylvanicus mainly in dense sedges and grasses, willows and a few on dry slopes but heavily vegetated areas bordering the marsh. Some used runways among the sedges & grasses. Noticed droppings along sedges and mosses lining Creek on lower reaches of creek but although good for *Microtus* now there were only an occasional one taken. Subadults in heavier more matted grasses of open fields.

Synaptomys borealis among sedges and grasses among willows. Soil wet and associated with *Microtus pennsylvanicus*.

Sorex palustris at edge of spring fed creek near base of boulders.

Clethrionomys gapperi, mainly on drier slopes of dense vegetation bordering the marsh area but also occasionally found in marsh.

Black duck in partial mact.

blue-winged teal in flock of 18

one female Common loon caught in a net set for fish. This bird had been seen the previous day fishing in the same area. This bird was prepared as a skeleton. 7 others seen flying over ^{head.}