



Bat net at Mill Pond (or 1/5 mi. S and 2/5 mi. W Lameshur Estate House)



This net placed at edge of trees bordering a muddy embayment and about 30' from the edge of the pond. It was placed for Noctilio but remarkable as it may seem, I did not catch a single bird or mammal in 23 days and night of continuous netting.

Bat net at Lameshur Estate House. Place across open veranda on east side of house.

Bat net at 1/10 mi. W Lameshur Estate House. net was stretched between ~~one~~ large poles



which I erected and a small tree on the side of the canyon so that the bottom of the net was 5 feet above a made-made, freshwater pond at the mouth of a canyon 1/10 of a mile

from the net at 1/5 mi. SW (or 1/5 mi. W.S.W) of Lameshur Estate House. The canyon supported large trees as far down as the pond and cut over clearings continued toward the beach. Each of 2 Stenoderma were in middle of net.

Sept. 4, 1957

LIST OF BIRDS NOTED AND COLLECTED.

NOTED

ON SAINT JOHN ISLAND AND ADJACENT OUTLIERS

<i>Pelicanus occidentalis</i>	<i>Sterna fuscata</i>	<i>Sericornis holoserena</i>
<i>Sula leucogaster</i>	<i>Sphrapicus varus</i> (holo)	<i>Orthorhynchus stiliis</i>
<i>Ardea herodias</i>	<i>Puffinus thomsoni</i>	<i>Tyrannus dominicensis</i>
<i>Buteo jamaicensis</i>		<i>Myiarchus stolidus</i>
<i>Falco sparverius</i>		<i>Margarops fuscatus</i>
<i>Haematopus ostralegus</i>		<i>Vireo altiloquus</i>
<i>Actitis macularia</i>		<i>Alcedo petechia</i>
<i>Columba leucocephalus</i>		<i>Leucurus aurocapillus</i>
<i>Columba squamasa</i>		<i>Coereba flaveola</i>
<i>Anthracoceros dominicus</i>		<i>Hirundo lunifrons</i>
<i>Elaenia martinica</i>		<i>Tyrannus bicolor</i>
<i>Progne dominicensis</i>		<i>Larus atricilla</i>
<i>Mimus polyglottis</i>		<i>Sterna albifrons</i>
<i>Casmerodius alba</i>		<i>Anous stolidus</i>
<i>Ardeola ibis</i>		<i>Ceryle alcyon</i>
<i>Nycticorax nycticorax</i>		<i>Crotophaga ani</i>

BIRDS COLLECTED