640823-87 annoyed by man. Singles fed in quet bays.
Junes hyemalis! State colored junes. Strigles fled among spruce & decedures forests at comp. Chordeles minor nighthaute. One or two fed ion lokes in evening, generally as singles Canachites Canadensis Sprice grouse noted one group only (see notes). These 8 fed in ground and flew to low brunches of trees. They gowld be approvehed to within 20 feet before flying. megaceryle aleyon. Ringsesher. they ded not call as frequently as in Kansas area. Litta Chuadensis. Red- breasted nuthatich Hard or seen regularly every day. warbling riveo (sp?) " One at camp on Lake III. Lang all morning up to 11:00 A.m and then stopped Inhabited Islend to west. Birds that will be affected by man and increase eise of area would melude: Common loon, red-breasted loon, block duck, Canada jay, roven, craw, gull, bacd eagle, Chipmunk, beover, painted tevilles and while-tailed deer. There are the kinds of animals that are attracted to comp sites or one annoyed by the presence or passing of man and Condes in the area, The common loon was being displaced continuously by passing boats as was the flack of young mergansers. Concentrated use of certain areas may preclude the nesting and feeding activities of the bold lagle. While beaver are mainly nactural, They are annoyed by consis. Painted tentles that spend Emsiduable time sunning on logs are always disladged by passing consertable those fishing near the edge of the labe. Fovortion of the block duck has produced a bird that is not at all like The wild duchs that are away from the trowelled routes. Chepments. Canada juy and crows has completely changed their normal feeding activities because of the soleculation of more for their presence around comps. Overfraking for

Inorthern puhe has produced a fish population that is

out of bulance with the natural population. notwal

populations of plants are being empletely changed at comp

sites. One week at a comp site will trompled ground cover

and remone dead foreword timber,