



extended directly back.
birds in the yard
in the course of the
meal, 2 bluejays, 1 wren.



the owl would straighten up to a perpendicular position and would remain in this position for as long as the source of intrusion was present. It would then return to its normal profile. When the suspecting a potential food source? it leans forward and drops head down.

at times it will move head back and forth. This action could develop into an action that involved the entire body in a circular -

back and forth movement; the body being raised and lowered, at some times to a propped position. In viewing food 6 inches below its perch, it will extend its head upward, draw chin in as far as possible, and attempt to fix vision on food. Another pose is to draw head into shoulder and ^{stare} upwards.

June 19, 1965

screech owl dove at food with claws first time today.

7/10 mi. E and 7/10 mi. N Clinton (P.O.), Douglas Co., Kansas

June 9, 1965

collected the following Bullocks Oriole & yellow-breasted Chat;

- 650609-1 Icteria galbula T.L. 197, wt. 38 gms ♂
- 650609-2 Icteria virens T.L. 175, wt. 26 gms ♂

In the afternoon while cruising in the area of the proposed Clinton Reservoir, recorded the following kinds of birds and mammals. rabbit (3 1/2 inches T.L.), marmot, fox squirrel, grey squirrel, cardinal, mourning dove, robin, Eng sparrow, cowbird, grackle, red-wing blackbird, turkey vulture, red-tailed hawk, titmouse, black capped chickadee, chimney swift, barn swallow, bank swallow, great blue heron, red bellied woodpecker, yellow shafted flicker, downy woodpecker, red headed woodpecker, pewee, phoebe, indigo bunting, Bullock oriole, orchard oriole, field sparrow, crested flycatcher, Eastern Kingbird, acadian flycatcher, blue jay, blue bird, ruby throated hummingbird (4), meadowlark, horned lark, summer tanager, goldfinch, killdeer, loggerhead shrike, brown thrasher, mockingbird, dickcissel, lark sparrow, crow, house wren, red-eyed vireo, yellow throat warbler,