

3 mi S divide, highway 287, Larimer Co., Wyoming.

and 2 does Aug 5, 1969

3 large buck (mule deer) and 5 antelope in same area just below camp. The does preceded the bucks. To west were 40 antelope. Others fed on distant ridges. Magpie & bluebirds dominant birds in morning. Left camp, mileage 28697 to Larimer Twp 68° F at 8:25. Sky 50% clear. Divide 28700. Pie Sidrug 28703. 12 antelope to W. From 28714 to outskirts (Cement Co) S of Larimer counted 14 *Citellus* along road right-of-way. Larimer 28721. At 5 mi. W Larimer noted a redtail hawk. Continued to Libby Lake at mileage 28

Set 140 traps in Krumholz NE of camp. Both prostrate and upright spruce on slopes. Soils bare and dry under Krumholz.

4/10 mi. S and 6/10 mi E of 10,850 ft.

Libby Lake, NW sec 17, R 79W, T 16N, Albany Co., Wyoming.

Aug 6, 1969

Collected the following mammals from trap line of 140 traps.

690806-1	<i>Clethrionomys</i>	124-39-18-11 ⁸ -15 gms testes 10 mm ♂
SK: 690806-2	<i>Peromyscus</i>	153-68- ²⁰ 18- ¹⁸ 20-16 gms ♂ testes 10 mm
690806-3	<i>Phenacomys</i>	113-28-18-12-14 gms. ♂ Testis 3 mm
690806-4	"	123-28-18-14-18 gms ♂ testes 3 mm

Birds in area: Canada jay, white Cr. sparrow, ^{gray headed} junco, pipit, mountain chickadee, Calif. gull, black tern, green tanager, water ouzel, pine siskin, water ouzel, black tern (?), robin.

Libby Flats, 10900 ft., Albany Co., Wyoming

Aug. 7, 1969

Last night set 140 traps (20' interval) among Krumholz on ridge just west of the quartzite rock field. This is the exact area trapped last year in August. Conditions are about the same as last year as concerns flora. This period of trapping is characterized by lack of moisture on plants in morning and windy days. The trapping area at Libby Lake has a different exposure and is not as well developed as the Libby Flat. There is more current shrubs at the edge of Krumholz at Libby Flat than at Libby Lake. Of the two areas, the Libby Flats is the more typical Krumholz.

