


Continued up trail. At the upper limits of before the benchland at base of peak, there is a noticeable terminal growth of krumholz across the S exposure of the mountain and a good place to set traps in the future, mainly because of its accessibility. In this area above the krumholz are rock exposures which show geometrical designs of yellow marks on the native rock. These marks are stains and show influence of border limits.

 Arrive at top of S exposure (at trail level) at altitude 11198. A hummingbird here (black tail & head, white rump on neck and gray body). From trail descended to W to edge of platform and beginning of krumholz. Photo 690814-14 of south edge showing influence of winds from the S-SW-W. (Controlling winds are from the NW). The lichens on rocks are characteristic of these upper rock fields. Photo 690814-15 of the lower edge of krumholz on W end of platform. Mt Reid in distance to N. These two above photographs at 11155 ft. The trapline of 140 traps started at this point at 11155 ft in foreground of 690814-15. Photo 690814-16 showing krumholz at the W end of platform where traps 1 to 70 were set (20' interval) in most favorable and sheltered parts of krumholz. This photo at 11242 ft elev. Tryon Lake Complex to W. Pampanagos Mt can be seen as faint outline above mesalike mountain in background. Photo 690814-17 at 11300 ft at upper limits of the massive krumholz on W end of platform. Shooting to W. From this point the krumholz is interrupted and rock fields and sparse soils separate it from the next and last group of krumholz to the east at base of peak (SE end platform). Traps across this rock field were set at 50' intervals. In the last group (highest on platform) krumholz trap set trap no. 121 to 138. From base of peak upper end platform and krumholz 11435 ft photographed round mountain to SE as no 690814-18. It appears that the underlying strata contribute to its profile as other similar masses can be seen beyond, Granddaddy Lobe Basin beyond. Arrive at base of peak at end krumholz at 6:20 P.M. At next switchback on peak to N set 2 traps in the most upper limits of krumholz on the S + SE exposure of Bald Mt, at 11527 feet. From this elevation to base of peak and on platform just beyond base is good area to trap for mammals that need rock protection and also some soil & vegetation. At ^{at} out 11600 were the greatest congregation of black rosy finch.