

Polymorphism for the occurrence of the second upper premolar in Felis bengalensis. Implications for the interpretation of fossil felid materials. Carnivore Genetics Newsletter 2(10):297-305. (With Neil B. Todd).

Gene frequencies of the domestic cat in Lawrence, Kansas. Carnivore Genetics Newsletter 3(1): (With Neil B. Todd).

Cat population genetics in the U. S. Southwest and Mexico. Carnivore Genetics Newsletter 3(1): (With Neil B. Todd and D. Creel).

Manuscripts in preparation

Microgeographic movements and asynchrony of seasonal population fluctuations in the cotton rat, Sigmodon hispidus Say and Ord 1825. (With Norman A. Slade) submitted Jour. Mamm.

Seasonal coexistence of Sigmodon hispidus and Microtus ochrogaster: The role of reproduction and population structure in competitive exclusion. (With Norman A. Slade).

Mammals of southern New Hampshire. (With Thomas H. Kunz and James G. Hullett).

Karen S. Harris

Spatial patterning in sympatric populations of Sciurus niger and S. carolinensis. 1976. Master's thesis, Univ. Kansas.

A three-dimensional home range model, ms submitted to J. Mamm. Koeppl, J. W., N. A. Slade, K. S. Harris, and Robert S. Hoffmann. The above is to be revised slightly and submitted to Ecology.

Lawrence R. Heaney

Manuscripts in press:

Distributional and natural history notes on some mammals from Pueblo, Mexico. Southwestern Naturalist (out this fall?)

Manuscripts in prep:

Island area and body size of insular mammals: evidence from the tri-colored squirrel (Callosciurus prevosti) of Southeast Asia.

Natural history of rock voles (Microtus chrotorrhinus) in Minnesota. (second author).

Glaucomys sabrinus. (Mammalian Species series account)

Limb proportions of the Holarctic ground squirrels (Marmotini, Sciuridae) (Junior author).

Checklist of the squirrels of the Oriental region (with J. W. Koeppl).