western sandpipers. once in a week
American Crow. Feed at various places within 3-4 miles range. May be absent in morning or afternoon but generally in area part of day. 18 is average size of flock.
dark-eyed junco. Most common bird of land birds and 40% of time at feeding station by Barth.
bald. eagle. Some days absent but 50% of time in area. generally in morning and then all morning.
western grebe. 40% of time.
downy woodpecker. Once per week.
Marbled murrelet. once every 4 days. Center bay
pigeon guillemot. once every 6 days. center bay
northwestern crow (identification questionable) but like American crow but 1/3 population.
herring gull. once every 3 days.
killdeer. once every 3 days.
Wilson snipe (common snipe) once every 12 days.
hooded merganser. once every 6 days. near shore
ruddy duck once every 10 days open waters
red-tailed hawk. once every 4 days.
Canvas-back once every 15 days.
ring-billed gull. once every 10 days.
mew gull once a day
savanna sparrow once every 5 days.
Thayer's gull. once every 10 days.
mallard. once every 15 days center bay
black turnstone once every 12 days.
bushtit once every 3 day
Chestnut-backed chickadee. once every 3 days
starling. once every 2 days.
golden-crowned kinglet. once every 3 days.
purple finch once every 12 days.
harbor seal. twice a day.
raccoon tracks mud in tidal zone new T.T.
black-tailed deer. Tracks uncommon.
ringed-billed gull. once every 4 days.
dunlin - once a week
western sandpiper - one 2 weeks.