clearing of negro-heads 4 feet high and thickly covering the ground. Two red-tail now at scene of sucking but no response from small birds

8:45 G.T. Towhee, 2 Aud. Warblers chipping in aspen.

8:55 Arrived at spring among aspen trees, being first water after leaving spring at camp this morning. Grass and willows line the springy area with small meadow ~~below~~ at base of hill where spring comes down. Two immature R. Hawk. Sapsuckers fed in willow. Approached with 8' before they finally flew. They were feeding upon the bark of the willow and remained close together. No adult in sight. The next spring visited beyond and on same trail is in a more cleared situation:



Failed to observe any mucrato runs but from general appearance am sure would support a few. The spring to the right is a mound type and is built up above the surrounding area. A few birch and not willow are found in center of area. Leaving spring and still trending to the north arrived at the Government constructed spring and trough in open flat below divide at 9:50. A robin, 4 G.T. Towhees, and 4 siskin were seen between last spring & Government spring. From the spring watched two golden eagles feeding along the west exposure of Timpanogos and about 1/5 the distant up mountain from our elevation. One flew just above the ground on hillside while the other bird flew higher and followed closely behind. The feeding procedure would be to fly south along side hill for about 5 blocks then return back to starting position at a higher elevation from surface of ground and then follow over same course about 15' above ground. I suspicion that the bird that followed and remained at a higher elevation was probably an