




immediate area in front Bee House out 4' to 12'. all otter captured candlefish. Continued fishing across entrance Spring Cove, remaining in close group, to Flat Point via cove, to π . where they put out for 25" then returned to water. Continued to Round Point, all but one put out, climbed to top of exposed rock, rested 20 seconds and then descended between Round and Flat points, thence fished in this cove and then put out on Flat Point to upper edge rocks to old sunning and drying area in grassy spot and after cleaning fur and marked territory left into cove between Flat and Dissected points and thence S. at times all seven almost contact, other times dispersed but within 10' diameter. Success in capturing food 80 per cent. whatever material used for food was chewed for from 10" to 60". No sign of aggressive action among otter. Diving from perfect or instantaneous to irregular. Saw no separation of adult and young interaction. The tide was high and 1' to 1 1/2' below flat part of Flat Point.

2:35 P.m. Common murre this

date



Townsend chipmunk has not been observed since last recorded in journal.

No mule deer up to at least 4:20 P.m.

5:25 Adult female mule deer and her fawn, 26" high at shoulders at Bee feeder.

5:30 P.m. Large mule deer with right eye line missing, fed at feeder then left and gained to cliffs and stopped. Young ♂ came down trail, past buck and went to feeder and fed for 3 minutes, then large buck came back to feeder and at 8' from young the young left area. After buck left feeder it stopped top of cliffs for 2-3 min, thence center of Deer Trail slope, thence along trail to π .

Ad. ♀ mule deer and her fawn and young ♀ came to feeder. For some reason the adult ♀ deer left at great speed followed by young ♀ and then fawn.

Deer (mule) come to feeder in winter later (light-wise) than