

oystercatchers, 41 mew gulls, 14 glaucous-winged gulls, 13 surf scoters, 8 buffleheads, 17 sanderlings, 12 American Crows, 1 great blue heron, 4 American widgeons,

4:10 P.M. In the sound between Leo Reef and Bee House. 18 mew gulls concentrated ^{feeding} in an area 10' x 6'. 4 or 5 gulls would rise ^{4'} above the group and then dive straight down into water, sometimes completely submerged for 2-3 seconds. If one was successful in capturing a fish, the other gulls would try to take the fish in flight. 3 double-crested cormorants swarm and fished among the gulls without interfering with their feeding behavior. 4 arctic loons fed at the periphery of the gull group. 3 red-necked grebe acted curious but did not participate in the feeding activity. These gulls perform this type of activity independent of association with other kinds of birds. They will, however, join groups of cormorants or other fish eating birds that operate in a group formation. 3 common murrelets fed approx. 150' away.

Observed an immature bald eagle in a tree at the S end of the causeway to Sperry Peninsula.

From above causeway to Humphrey Head counted 81 American robins. No groups of American Crows.

At Bee House, no mule deer yesterday of previous night but this evening a young ♂ + ♀ came to feeder after dark.

Prepared a specimen this evening from Bee House.

at Humphrey Head, Lopez Island, San Juan Co., Washington:

870220-1 *Peromyscus maniculatus hollisteri* 178-80-23-19-28 gms. ♂, testes 8 mm diameter skinned tail 10 mm from base is 2.3 mm.

Feb 21, 1987

Two bald eagle sighting between 7:00-8:00 A.M. Sky overcast and water calm.

1:00 P.M. 18 harbor seal on Leo Reef.

Daughter Mary and I followed beach from Odlin Park to Flat Point. she continued S to Lopez Village along the beach, I returned to Odlin Park and then drove to Village to wait for Mary. For the 1 1/2 mi. distance to Flat Point (between 2:00 P.M. to 4:30 P.M.) made the following observations: (numbers are totals for the one way trip to Flat Point) 28 glaucous-winged gulls, 53 buffleheads, 13 greater scaup, 1 immature bald eagle, 6 pelagic cormorants. The gadwall (*Penopus generosa*) are common, on the last 2/10 miles of trip as we