

It was successful in displacing 80% of the incoming birds of which at this time was 1-3. On two occasions, one a male and the other a female, the sick bird was driven from the feeder. and in all cases to lower branches of trees 15-20' away. In one of the case the female continued its harassment and drove the sick birds 150' into Spring Cove. ^(sick one) It returned 10 minutes later. The flight is slow and direct to wherever it goes, not the erratic and energetic flight of the normal birds. Eyes are frequently closed for a second or so. I would say the bird has slightly improved in activity this morning compared to yesterday evening.

9:10 A.M. Imm. bald eagle flew NW 10' above water and landed on high tide rocks on Lichen Point. It hopped to high ground and remained for 4 minutes, then left to S. Basal hill light only.

10:30 A.M. sick hummingbird respiration 126 per minute.

10:35 A.M. Algae (mainly greenish ell grass) along shore to 50' from Boundary Point to Lichen Point, at least, and moving S with currents.

10:40 Ad bald eagle and 2 immature bald eagles in trees above Flate Point and Dissected Point. One of the immatures had a scrap of food & adult attempted to get the food from the imm. After 3 attempts (among limbs of Douglas fir) it flew to Lichen Point and thence E to Blahely at point above New Willow Island. At midway it dropped from 300' to 75' in 45° angle but changed course and again gained 300' to Blahely. It then flew N along ^W side Blahely at 1/3 height of mt. I lost it at Thatcher. But in about the same time it took the bird to fly to Blahely it returned (could have been another adult) and again attempted to take away a new scrape of food (4"x6") that one of the young had just picked from the water. It chased the immature into Spring Cove, thence to 10' NE of Tide Point where it attempted, unsuccessfully, to get the food from the immature. Both birds then flew to tree (Douglas fir) above Dissected Point where the immature dropped down to the hillside thru a tangle of bows. The adult did not pursue the immature which completely consumed the food on the ground. Both birds then left to S leaving the 3rd immature who had remained on a dead limb 30' from activity for 25 minutes.