

a large ad ♀, dominant and aggressive. She tolerates others for 2-3 seconds, then thrust front feet or pushes away with head or neck and if this does not displace other deer, she will aggressively and forcefully charge or even take upright stance, in other words she expresses mild aggressive and then gets mad and continues with force.

Sister, a mule deer, 2nd year. } pair of mamma Deer's fawns,  
Buster, a mule deer, 3rd year }

A single fawn ♂ (brother of twin ♀) of mamma Deer.

Ad. female and single fawn. Competitive size physical condition good but the least aggressive and <sup>is</sup> timid

Total 3 adults, 2 subadults, 4 fawns.

General observations:

(an orphan)

All adults and subadults chase juvenile ♂ of mamma Deer most deer, especially fawn come into feeding area with tails held 90°-120° then drop tail to body at feeder

Twins of light color ♀, groomed each other at feeder. Elsewhere the mother + two twins groomed each other at 75' from feeder.

Buster, in his own rank, disperses fawns.

One of the twins <sup>of light colored ♀</sup> attempted mounted the mother. On several occasions when one of the twins would touch the <sup>top of</sup> rump of the ♀, she (the ad) would crouch down for a second or so.

young fawn to presence of ad ♂, especially as they leave the feeding area if displaced by the ♂.

8:30 A.M. <sup>a single</sup> harbor seal, 40' from shore, done for 2 minutes and 20 seconds.

9:15 A.M. Single river otter swam NE past Bee House, 20' from shore to Boundary Point and beyond. At one time it remained underwater 38 secs. 8-10 sec is usual surface time when otter are moving forward and hunting at the same time.

9:20 A.M. 1 western grebe out 25' from shore below Bee House.

Between approx 8:30 A.M. to 11:00 A.M. an orange of 9 Bonaparte gulls hunted waters of Spring Cove Bay (area between Lichen Point and Boundary Point). They were from water contact to 20' high most of the time.

12:30 P.M. western grebe fishing for candlefish 2'-6' from shore directly below Bee House. Above successfully 3 times, each time coming up from under fish and causing them to scatter on the surface. Most of the time the bird made 2-3 continuous attempts within the surface fish before making a catch. A Bonaparte gull capture one of the candlefish when fish were