


8:30 A.M. The top-eared ad. ♀, 31" at shoulders at feeder at Bee House, also ♀, 30" at shoulders at Bee feeder. Her two fawns remained on slope above house and did not come to feeder. They instinctively are shy of man and will run up hill and take to ground cover. An antlered (1" long) ♂ touched nose with above female and was comfortable at feeder with the ad. ♀. The summer pelage (golden brown) completed to neck in front of shoulders. The old winter pelage ^{on neck} still retains spots of a previous juvenal pelage.

Trip to Anacortes & Mt. Vernon. Tidal marks on E side Flower Island distinct.

Big Flower Lick is eroded flat on top at ^{top} barnacle zone.





at W end Thatcher Pass, 3 pigeon guillemot.

Pointer Island usual glaucous-winged gulls on territories. (territories 3 times closer than those on Flower Island.) and 11 cormorants resting on top of E end island.

2 glaucous-winged gulls, 2 glaucous-winged gulls, 3 pigeon guillemots. Arrived Anacortes.

Continued thru Anacortes and drove clockwise around March Peninsula. On NE point of March Peninsula for 1 mi. shoreline 42 great blue heron in groups of 6, 6, 5, 14, 4, 7.

On W side March Peninsula water approx 3 acres and 1,100 drainage 5 of causeway (R, R) . Glaucous-winged gulls average 1 per 200' along outer edge  of tideland. No brant in Padilla Bay.

At La Conner at Chandler marine store, the owner reported a coot arrived 3 days ago and has remained in area. Ordinarily coot are fresh water birds. He also reports that all winter birds have left the area.

Continued to Thousand Trails after checking with Mr. Johnson in Mt. Vernon. At 9:30 P.M. high tide at Lone Tree Point. High water and waves come across ^{entire} connection between Lone Tree Point and mainland carrying large logs ⁽³⁰⁾ and waves. The gravel spit on N side lagoon covered with water from base of peninsula island to W side ^{101 west} Indian Lodge. Outlet channel now water back