


♀ mule deer 28" high at shoulders:
winter pelage showing in summer
golden-brown pelage.



Aug. 20, 1989

Harbor seal:

Leo Reef, rocks 3-6 are 16 seal

Little Flower Rock is 1 seal

North Flower Rock are 9 seal


Total 26

10:05 A.M. To Bee Boat moored in S end Shoal Bay. Rain last night. Belted kingfisher on top mast, flew to mainland. From boat caught a Leptocestus armatus, Pacific stag-horn sculpin. It measured 230 mm standard length and weighed 255 grams.

Aug. 21, 1989

95% fog and overcast.

♂ mule deer, 2 points, ^{no} velvet and dominant ♂. Some summer pelage still between ears and antlers.



♀ mule deer first to molt and is the most active and unconfiding of ♀'s.

♂ molted, antlers clean of velvet.

♂ unmolted, 33" at shoulders and antlers free of velvet.

♂ " and velvet still on antlers. Above 3 at feeder at same time. Molt starts from head, ^{and} moves posteriorly to ears and antler areas of head.

Photo 890821-1 of wind sailing in Spring Cove. One of O'Leary's renters.

2 bats started evening flight at 8:24 P.M. and continued to dark at least.

Aug. 22, 1989

Buster, a ♀ mule deer had 2 fawns, 1 23" high at shoulders ♂ with antler buds under the skin. The second one 22" high at shoulders. Both have a new dark dorsal median line of new pelage but both with spots in pelage.