


13 pleasure craft visible at one time, 90% going E. Only 2 glaucous-winged birds observed on entire trip. Returned to Lopez Island approx. 7:00 P.M. 2 pigeon guillemots, 7 pigeon guillemots, 2 great blue herons flying 100' high in middle Rosario Straits, they flew to James Island then S. 2 glaucous-winged gulls w Thatcher Pass. Continued to Lopez Island.

May 27, 1991

Deer at Bee House feeder: mule deer.

δ 25" high at shoulders, fawn this year 18" high at shoulders, mother of fawn 30" high at shoulders, ^(Year) male with antlers 1" high, ♀ 27" high at shoulders, $\frac{1}{2}$ molted.

σ 33" high at shoulders →



May 28, 1991

Leo Rocks 4 to 7, 30 harbor seal.

" " 9 , 20 " "

Little Flower Rock, 9 " "

Flower Island, 9 glaucous-winged gulls,

At bird feeder, American Crow and eastern gray squirrel 6" apart.
no boats at Spencer Spit, because of cold weather.

May 29, 1991

910529-1, ^(im.) Juncos hyemalis ^{mm} 151 TL, 73 wing, 59 tail, 22 tarsus, 9.5 exposed culmen, 340 wing spread, 18 gms wt. Measurements only.

2:20 P.M. River otter swam N past Bee House. It fed within 1' from the shore all along the way. Bald eagle on nest SE Frost Island, 27 glaucous-winged gulls Flower Island. No harbor seal in area.

^(mule deer) fawn this year 15" high at shoulders, mother left front leg injured, let fawn nurse. This fawn has white hair in tarsal leg gland, scratches ear with hind foot, licked tail area and after 1 min. difficult to use hind leg, bunts mother's head, frolics with and around mother, traverses rock slopes, touched tail of eastern gray squirrel and jumped back, reacts to squirrel, Townsend Chipmunk, Rufous-sided Towhee more than adults of the mule deer, other deer accepts fawn.

σ 26" high at shoulders $\frac{1}{3}$ molted on sides.
most all deer show new summer hair
thru winter pelage on sides →
Large bucks complete body pelage (summer)
Some large ♀ just showing new hair.

