ÆLURŒDUS VIRIDIS, *(Lath.)*

W. Hart del. et lith.

Mintern Bros. imp.

ÆLU

Green Grakle, Lath. Gen. S
Gracula viridis, Lath. Ind.
Lanius crassirostris, Paykul
Kitta virescens, Temm. Pl. C
Ptilonorhynchus smithii, Vig
p. 325 (1846).—
Ptilorhynchus smithii, Bp. C
Ailurœdus smithii, Cab. Mu
Ælurœdus crassirostris, Scla
Linn. Soc. N. S.
Ptilonorhynchus crassirostri
Ælurœdus viridis, Sharpe, C
North, Nests &

In nearly every country in
bestowed upon it. In Nor
and in Australia the presen
resemblance of its note to t
Bower-birds, but it has new
in which this bird is found,
respecting its habits, as I c
Satin-bird; but I could n
which it breeds; it is doub
As will be seen below, the
never heard of any "bower

The range of the Cat-bir
Queensland. He writes:—
Illawarra district, and is fo
becoming scarcer, however,
hood of the Clarence, Richm
but not so freely dispersed,
knowledge extends, this spe
that extend along the easte
the Macleay, and the Clare
localities in which it may b
to the habits of the Cat-bird
tree."

It should be noted that
present species; but I pres
birds, as Mr. North does no

Mr. Gould gives the foll
cherry, when in season, affo
finding any remains in the s
a wary bird, little caution b
the lofty branches of the tre
note which differs so much
mistaken. In comparing it
my readers a more perfect i
concert is performed either