

SHOOTING JARKE

Shooting Jarke, 1900

Shooting Jarke, 1900

This species, the white-bellied, is distributed in the north-western and eastern parts of the island, and is the most numerous of the group. It is a very common bird, and is often seen in the open country. The male is much larger than the female, and has a much longer bill. The female is much smaller, and has a much shorter bill. The male is much more brightly colored than the female, and has a much more extensive white patch on the throat. The female is much more drab, and has a much more extensive brown patch on the throat. The male is much more active than the female, and has a much more extensive white patch on the throat. The female is much more sedate, and has a much more extensive brown patch on the throat.

The male is much more brightly colored than the female, and has a much more extensive white patch on the throat. The female is much more drab, and has a much more extensive brown patch on the throat.

The male is much more active than the female, and has a much more extensive white patch on the throat. The female is much more sedate, and has a much more extensive brown patch on the throat.

The male is much more brightly colored than the female, and has a much more extensive white patch on the throat. The female is much more drab, and has a much more extensive brown patch on the throat.

The male is much more active than the female, and has a much more extensive white patch on the throat. The female is much more sedate, and has a much more extensive brown patch on the throat.

The male is much more brightly colored than the female, and has a much more extensive white patch on the throat. The female is much more drab, and has a much more extensive brown patch on the throat.