## CHARADRIUS VEREDUS, Gould.

T. MAN. W. TO. A. O. D. W. M. M. W. W. C. A. O. W. C.

Brown Plover.

Charadrius veredus, Gould in Proc. of Zool. Soc., February 22, 1848.

A specimen of this species was procured at Port Essington by Mr. Gilbert, and a second has been sent to me from Sydney by Mr. Strange; these, and one other example, are all that have come under my observation; we may consequently infer that it is a very rare bird in Australia. Judging from its structure and the character of its plumage, it would seem to be nearly allied to the restricted genus *Eudromias* or Dottrells: I have however placed it in the genus *Charadrius*, until a knowledge of its habits, whether or not it undergoes any periodical changes of plumage, and if the specimens from which I have taken my figures be perfectly adult, have been ascertained,

Crown of the head and all the upper surface brown, each feather narrowly fringed with buff; primaries blackish brown, the shaft of the first white; tail brown, narrowly edged with white, the brown colour gradually fading as the feathers recede from the centre; face, a broad stripe over the eye and the chin buffy white; sides and back of the neck and the breast buffy brown; abdomen and under surface white; irides very dark brown; legs and feet brownish flesh-colour; bill dark brown.

TY, M to W at M. outhor M

The figures are of the natural size.