

19.

SELEUCIDES NIGRICANS (*Shaw*).

Twelve-wired Bird of Paradise.

- Oiseau de Paradis blanc*, Forster, in Forrest, Voy. Moluq. et Nouv. Guin. p. 160 (1780: ex Valenty).
Oiseau de Paradis noir et peu connu, Forst. t. c. pp. 154, 160. no. 6 (1780: ex Valenty).
Paradisæ candida (pt.), Forst. Zool. Ind. pp. 31, 35, 36, var. secunda (1781: ex Valenty).
Paradisæ ignota, Forst. t. c. pp. 31, 36 (1781: ex Valenty).
Paradisæ alba, var. β , Gm. Syst. Nat. i. p. 402 (1788: ex Forst.).—Gerini, Ornith. tab. 65. fig. 1.—Lath. Ind. Orn. i. p. 197 (1790).—Blumenb. Abbild. naturh. Gegen. pl. 96 (1810).—Bechst. Kurze Uebers. p. 133 (1811).—Cuv. Règn. Anim. i. p. 403 (1817).—Schl. J. f. O. 1861, p. 386.
Paradisæ melanoleuca, Daud. Tr. d'Orn. ii. p. 278 (1800).
Le Manucode à douze filets, Vieill. Ois. Dor. ii. p. 29, pl. 13 (1802).
Le Nébuleux, Levaill. Ois. Parad. i. pls. 16, 17 (1806).
Paradisæ nigricans, Shaw, Gen. Zool. vii. pt. 2, p. 489, pls. 60, 61 (1809).—Steph. Gen. Zool. xiv. pt. 1, p. 76 (1826).
Paradisæ violacea, Bechst. Kurze Uebers. p. 133 (1811).
Paradisæ vaillanti, Shaw, Nat. Misc. xxiv. pl. 1025.
Le Promerops multifil, Levaill. Hist. Nat. Prom. et Guép. p. 38, pl. 17 (1806).—Less. Man. d'Orn. ii. p. 6 (1828).
Falcinellus resplendescens, Vieill. N. Dict. d'Hist. Nat. xxviii. p. 165 (1819).—Id. Enc. Méth. p. 581 (1823).—Id. Gal. Ois. i. p. 307, pl. 185 (1825).—Less. Traité d'Orn. p. 322 (1831).—Drap. Dict. Class. xiv. p. 293 (1845).
Epimachus albus, Temm. Man. d'Orn. i. Introd. p. lxxxvi (1820).—Wagl. Syst. Av., *Epimachus*, sp. 9, p. 121 (1827).—Cuv. Règne Anim. i. p. 440 (1829).—Gray, Gen. B. ii. p. 94 (1848).—Id. P. Z. S. 1858, p. 190.—Id. Cat. B. New Guin. pp. 21, 55 (1859).—Id. P. Z. S. 1861, p. 435.—Wallace, P. Z. S. 1862, p. 160.—Finsch, Neu-Guinea, p. 165 (1865).—Schl. Mus. Pays-Bas, Coraces, p. 95 (1867).—Id. Nederl. Tijdschr. Dierk. iv. p. 49 (1871).—Musschenbr. Dagboek, pp. 202, 234 (1883).—Rosenb. Mitth. orn. Ver. Wien, 1885, p. 53.
Twelve-wired Paradise-bird, Lath. Gen. Hist. iii. p. 199, pl. 48 (1822).
Seleucides acanthylis, Less. Ois. Parad. Syn. p. 29, Hist. Nat. p. 229, pls. 36, 37 (σ ad.), pl. 38 (σ juv.) (1835).—Id. Compl. de Buff., Ois. p. 540 (1838).
Nematophora alba, Gray, List Gen. B. p. 12 (1840).
Seleucides albus, Gray, List Gen. B. Addenda, p. 1 (1840).—Id. List Gen. 2nd ed. p. 15 (1841).—Bp. Consp. Av. i. p. 412 (1850).—Cab. Mus. Hein. i. p. 215 (1851).—Scl. Journ. Linn. Soc. ii. p. 163 (1858).—Gray, Hand-l. B. i. p. 105, no. 1275 (1869).—D'Alb. Ann. Mus. Civ. Gen. x. pp. 13, 20 (1877).—Id. Nuova Guin. p. 484 (1880).
Seleucides alba, Reichenb. Scansoriæ, p. 331, tab. 612. figs. 4092-93 (1853).—Rosenb. J. f. O. 1861, p. 46.—Wallace, P. Z. S. 1862, pp. 155, 158, 160.—Id. Malay Archip. ii. pp. 388, 412, 419, cum tab. (1869).—Elliot, Monogr. Parad. pl. xxii. (1873).—Scl. P. Z. S. 1873, p. 697.—Beccari, Ann. Mus. Civ. Gen. vii. p. 713 (1875).—Salvad. t. c. p. 785.—D'Alb. Nuova Guin. pp. 34, 345, 372, 456, 484, 485 (1880).—Eudes-Deslongch. Ann. Mus. d'Hist. Nat. Caen, i. p. 13 (1880).—Scl. P. Z. S. 1881, p. 450.—D'Hamonv. Bull. Soc. Zool. France, xi. p. 509 (1886).
Ptiloris nebulosus, Licht. Nomencl. Av. Mus. Berol. p. 10 (1854).
Seleucides resplendens, Rosenb. Naturk. Tijdschr. Nederl. Ind. xxv. p. 238 (1863).—Id. J. f. O. 1864, p. 123.
Epimachus resplendens, Rosenb. Reist. naar Geelvinkb. pp. 101, 116 (1875).
Seleucides ignota, Salvad. Ann. Mus. Civ. Gen. viii. p. 403 (1876), ix. p. 191 (1876), x. p. 154 (1877).—D'Alb. & Salvad. op. cit. xiv. p. 107 (1879).—D'Alb. Nuova Guin. pp. 582, 588 (1880).
Seleucides niger (nec Shaw), Sharpe, Cat. B. Brit. Mus. iii. p. 159 (1877).—Forbes, P. Z. S. 1882, p. 333.—Goodwin, Ibis, 1890, p. 150.
Epimachus resplendens, Rosenb. Malay. Archip. p. 552 (1879).
Seleucides nigricans, Salvad. Orn. Papuasias, etc. ii. pp. 561, 680 (1881).—Gould's B. New Guinea, i. pls. 14, 15 (1881).—Guillem. P. Z. S. 1885, p. 650 (Salawati).—Salvad. Aggiunte Orn. Papuasias, etc. ii. p. 157 (1890), iii. p. 241 (1891).—Sharpe, Bull. Brit. Orn. Club, iv. p. xii (1894).—Salvad. Ann. Mus. Civ. Gen. (2) xvi. p. 109 (1896).—Rothschild, Novit. Zool. iii. pp. 531, 633 (1896).

As Count Salvadori has pointed out, the earlier names of *Paradisæ alba*, *P. ignota*, and *P. melanoleuca* are not applicable to this bird, as all of them convey a false impression of its coloration, or are founded