PTILORIS PARADISEUS, Swains.

Rifle Bird.

Ptiloris paradiseus, Swains. Zool. Journ., vol. i. p. 481.—Ib. Class. of Birds, vol. ii. p. 331.—Jard. and Selb. Ill. Orn., vol. i. pls. 43 male, 44 female.—Steph. Cont. of Shaw's Gen. Zool., vol. xiv. p. 267.

Ptiloris paradisea, G. R. Gray, List of Gen. of Birds, 2nd edit., p. 15.

Hitherto this magnificent bird has only been discovered in the brushes of the south-eastern portion of Australia; so limited in fact does its range of habitat seem to be, that the river Hunter to the southward, and Moreton Bay to the eastward, may be considered its natural boundaries in either direction. I have been informed by several persons who have seen it in its native wilds that it possesses many habits in common with the Climacteri, and that it ascends the upright boles of trees precisely after the manner of those birds. It was a source of regret to me that I had no opportunity of verifying these assertions, but an examination of the structure of the bird induces me to believe that it is more nearly allied to the members of that genus than to any other: that its powers of flight are very limited, is certain from the shortness and peculiarly truncate form of the wing, and this mode of progression is doubtless seldom resorted to further than to transport the bird from tree to tree, or from one part of the forest to another. That it is stationary in the brushes of the Clarence, MacLeay, and all similar districts between the river Hunter and the settlement at Moreton Bay, is evident from the numerous specimens of all ages that have been sent from thence to Europe.

The Ptiloris paradiseus is without exception the most gorgeously plumaged bird yet discovered in Australia: the sexes offer the greatest possible difference in the colouring of their plumage; for while the male is adorned with hues only equalled by some species of the Trochilidæ or Humming-Birds, the dress of the female is as sombre as can well be imagined. The law which exists wherever there is a great difference in the colouring of the sexes causes many parti-coloured changes of plumage in the immature males during the period of moulting, which however during the first year, and probably for a longer period, cannot be distinguished from the females.

The adult male has the general plumage rich velvety black, glossed on the upper surface with brownish lilac; under surface similar to the upper, but all the feathers of the abdomen and flanks broadly margined with rich olive-green; feathers of the head and throat small, scale-like, and of a shining metallic blue-green; two centre tail-feathers rich shining metallic green, the remainder deep black; bill and feet black.

The female has the whole of the upper surface greyish brown; the wings and tail edged with ferruginous; the feathers of the head with a narrow line of white down the centre; line passing down the side of the head from behind the eye, chin and throat, buffy white; all the under surface deep buff, each feather with a black arrow-headed-shaped mark near the tip.

The Plate represents two males and a female of the size of life.