

n, vii. no. 2,

he Dresden
chenbroek's
a made up

—Rothschild,

chen. J. f. O.
onogr. Parad.

Dresden, vii.

, Parad. p. 43
895).

. xiii (1894).—

sden, vii. no. 2,

. Range, in
the feet greyish
a.' The adult
greyish (horn-

GENUS XVII. LAMPROTHORAX.

Add:—

Lamprothorax wilhelminæ, Sharpe, Monogr. Parad. part vi. (1896).—Meyer, Abhandl. k. zool. Mus. Dresden, vii. no. 2, p. 51 (1898).—Rothschild, Thierreich, Lief. 2, Parad. p. 21 (1898).

GENUS XVIII. CICINNURUS.

Add:—

Paradisea regia, Marshall, Arch. Néerl. vi. p. 297 (1871).
Cicinnurus regius, Salvad. Ann. Mus. Genov. (2) x. p. 822 (1891).—Meyer, Abhandl. k. zool. Mus. Dresden, 1891, p. 12; 1893, p. 23.—Sharpe, Monogr. Parad. part i. (1891).—Id. Bull. Brit. Orn. Club, iv. p. xiii (1894).—Madarász, Aquila, i. p. 91 (1894).—Salvad. Ann. Mus. Genov. (2) xvi. p. 822 (1896).—Reichen. J. f. O. 1897, p. 214.—Meyer, Abhandl. k. zool. Mus. Dresden, vii. no. 2, p. 47 (1898).
Cicinnurus regius (typicus), Rothschild, Thierreich, Lief. 2, Parad. p. 35 (1898).

Dr. Loria obtained a large series of specimens at Bara-Bara in Milne Bay, and at other villages on the Kemp Welch River to the east of Port Moresby. The iris was dull chestnut, the bill yellow, and the feet blue. The native name of the female bird was "Uanaro Uapu" (*cf.* Salvad. *l. c.*). Procured by Fenichel in the Finisterre Mountains, and by Tappenbeck on the Nuru River in German New Guinea.

Dr. A. B. Meyer (*l. c.*) says that the species was met with commonly by him during his journey across North-western New Guinea. It lives on the high trees, and he says that the blue colour of the feet is lovely, the claws grey, the bill clear yellow, and the iris clear brown. He also noticed the change of the colour from brown to red without a moult.

Add:—

Cicinnurus regius coccineifrons, Meyer, Abhandl. k. zool. Mus. Dresden, vii. no. 2 (1898).—Rothschild, Thierreich, Lief. 2, Parad. p. 35 (1898).
Cicinnurus coccineifrons, Sharpe, Monogr. Parad. part viii. (1898).

GENUS XIX. RHIPIDORNIS.

Add:—

1. RHIPIDORNIS GULIELMI TERTII.

Rhipidornis gulielmi tertii, Sharpe, Monogr. Parad. part ii. (1893).—Id. Bull. Brit. Orn. Club, iv. p. xiii (1894).
Rhipidornis gulielmertii, Meyer, Abhandl. k. zool. Mus. Dresden, vii. no. 2, p. 48 (1898).
Diphylloides gulielmertii, Rothschild, Thierreich, Lief. 2, Parad. p. 36 (1898).

GENUS XX. DIPHYLLODES.

Add:—

1. DIPHYLLODES MAGNIFICA.

Diphylloides magnifica, Sharpe, Bull. Brit. Orn. Club, iv. p. xiii (1894).—Id. Monogr. Parad. part vi. (1896).—Salvad. Ann. Mus. Civic. Genov. (2) xvi. p. 111 (1896).—Meyer, Abhandl. k. zool. Mus. Dresden, vii. no. 2, p. 48 (1898).—Rothschild, Novit. Zool. v. p. 87 (1898).
Diphylloides magnificus, Rothschild, Thierreich, Lief. 2, Parad. p. 36 (1898).

Dr. Meyer gives the following note from his diary of 1873:—"Its cry is a piping 'krrrrr, krrrrr.' Iris and feet black. Frequents the ground. The male clears a large round place: during this operation the female sits close at hand on the lianas or rattans, and when the male has finished, she dances with him on the newly-cleared space."