

Key to the Species.

- a.* Outer aspect of wings more or less purple.
a'. Under surface of body metallic steel-green or purple; feathers of the crown and throat steel-blue or metallic green.
a''. Head not in strong contrast to the back.
a³. Head dark steel-blue like the back
b³. Head metallic green, slightly contrasting with the steel-blue of the back.
b''. Head steel-blue; back purple like the wings
b'. Under surface of body black, with scarcely any metallic gloss; feathers of head and throat metallic oily-green
b. Outer aspect of quills always green and not purplish; general colour above and below metallic green

XXV. MANUCODIA.

- Manucodia*, Bodd. Tabl. Pl. Enl. p. 39 (1788)
Range. New Guinea, Salawati, Jobi, Mysol, Waigiou, Ghemien, Batanta, Aru Islands.

Key to the Species.

- a.* Intercapillary region rich purple, crinkled with transverse bands of velvety black
b. Intercapillary region metallic steel-green with purple reflections, but without recurved velvety margins to the feathers

XXVI. EUCORAX.

- Eucorax*, Sharpe, Bull. Brit. Orn. Club, iv. p. xv (1894)
Range. D'Entrecasteaux Islands.

XXVII. LYCOCORAX.

- Lycocorax*, Bp. Comptes Rendus, xxxvii. p. 829 (1853)
Range. Batchian, Halmahéra, Morotai, Raou, Obi Islands.

Key to the Species.

- a.* Primaries ashy brown below, somewhat paler, but not white, at the base of the inner web.
a'. General colour black, slightly washed with green; secondaries brown
b'. General colour dark rifle-green; secondaries black, washed with green
b. Primaries white for the basal half of the inner web

Type.

M. chalybeata.1. *chalybeata* and allies.2. *atra*.

Type.

E. corvii.

Type.

L. pyrrhopterus.

1. *pyrrhopterus*.
2. *obiensis*.
3. *morotensis*.

XXVIII. PARADIGALLA.

- Paradigalla*, Less. Ois. Parad. 1835, p. 242
Range. New Guinea.

Type.

P. carunculata.