

INTRODUCTION.

XVIII. CICINNURUS.

- Cicinnurus*, Vieill. N. Dict. d'Hist. Nat. xix. p. 247 (1818) *C. regius*.
Range. New Guinea, Salawati, Jobi, Mysol, Arn Islands.

Key to the Species.

- a.* Crimson above; the head like the back, with the forehead inclining to orange 1. *regius*.
b. Crimson above, but the forehead not orange, but crimson like the crown 2. *coccineifrons*.

XIX. RHIPIDORNIS.

- Rhipidornis*, Salvad. Ann. Mus. Civic. Genov. ix. p. 192 (1876) Type.
R. gulielmi tertii.
Range. New Guinea.

XX. DIPHYLLODES.

- Diphyllodes*, Less. Ois. Parad. Syn. p. 5 (1835) Type.
D. magnifica.
Cricocercus, Gloger, Handb. Naturg. p. 345 (1842) *D. magnifica*.
Range. New Guinea, Salawati, Jobi, Mysol, Waigiau.

Key to the Species.

- a.* Head dull brownish, slightly tawny on the sides of the hinder crown; upper back deep claret-colour.
d. Secondaries clay-coloured 1. *magnifica*.
b. Secondaries ochraceous orange 2. *seleucides*.
c. Secondaries golden orange 3. *chrysoptera*.
b. Head reddish-brown; sides of hinder crown conspicuously more rufous; upper back deep crimson.
d. Secondaries golden yellow 4. *xanthoptera*.
e. Secondaries golden orange 5. *hunsteini*.

XXI. SCHLEGELIA.

- Schlegelia*, Bernst. Ned. Tijdschr. Dierk. ii. p. 1 (1864) Type.
S. respublica.
Range. Waigiau and Batanta.

XXII. PTERIDOPHORA.

- Pteridophora*, Meyer, Bull. Brit. Orn. Club, iv. p. xi (1894) Type.
P. alberti.
Range. N.W. New Guinea.

XXIII. SEMIOPTERA.

- Semioptera*, Gray, P. Z. S. 1859, p. 130 Type.
S. wallacii.
Range. Batchian and Halmahéra.

XXIV. PHONYGAMA.

- Phonygama*, Less. Voy. Coquille, i. p. 636 (1826) Type.
Chalybeus, Cuvier, Règne Anim. i. p. 354 (1829) *P. keraudreni*.
P. keraudreni.
Range. Papuan Islands, D'Entrecasteaux Islands, N. Australia.